

AMALGAMATED FREE PRODUCTS OF w -RIGID FACTORS AND CALCULATION OF THEIR SYMMETRY GROUPS

A. IOANA, J. PETERSON AND S. POPA

University of California, Los Angeles

ABSTRACT. We consider amalgamated free product II_1 factors $M = M_1 *_B M_2 *_B \dots$ and use “deformation/rigidity” and “intertwining” techniques to prove that any relatively rigid von Neumann subalgebra $Q \subset M$ can be unitary conjugated into one of the M_i ’s. We apply this to the case M_i are w -rigid II_1 factors, with B equal to either $\mathbb{C}$, to a Cartan subalgebra A in M_i , or to a regular hyperfinite II_1 subfactor R in M_i , to obtain the following type of unique decomposition results, à la Bass-Serre: If $M = (N_1 *_C N_2 *_C \dots)^t$, for some $t > 0$ and some other similar inclusions of algebras $C \subset N_j$ then, after a permutation of indices, $(B \subset M_i)$ is inner conjugate to $(C \subset N_i)^t$, $\forall i$. Taking $B = \mathbb{C}$ and $M_i = (L(\mathbb{Z}^2 \rtimes \mathbb{F}_2))^{t_i}$, with $\{t_i\}_{i \geq 1} = S$ a given countable subgroup of $\mathbb{R}_+^*$, we obtain continuously many non stably isomorphic factors M with fundamental group $\mathcal{F}(M)$ equal to S . For $B = A$, we obtain a new class of factors M with unique Cartan subalgebra decomposition, with a large subclass satisfying $\mathcal{F}(M) = \{1\}$ and $\text{Out}(M)$ abelian and calculable. Taking $B = R$, we get examples of factors with $\mathcal{F}(M) = \{1\}$, $\text{Out}(M) = K$, for any given separable compact abelian group K .

Content

0. Introduction
 1. Conjugating subalgebras in AFP factors
 2. Deformations of AFP factors
 3. Deformation/rigidity arguments
 4. Existence of intertwining bimodules
 5. Rigid subalgebras in AFP factors: General Bass-Serre type results
 6. Amalgamation over $\mathbb{C}$: Free product factors with prescribed $\mathcal{F}(M)$
 7. Amalgamation over Cartan subalgebras: vNE/OE rigidity results
 8. Amalgamation over R : Factors with no outer automorphisms
- Appendix: Constructing freely independent actions

J.P. supported in part by NSF VIGRE Grant DMS 9983726 at UCLA; S.P. supported in part by nsf-grant 0100883.

0. INTRODUCTION.

We prove in this paper a series of rigidity results for amalgamated free product (hereafter abbreviated AFP) II_1 factors $M = M_1 *_B M_2$ which can be viewed as von Neumann algebra versions of the “subgroup theorems” and “isomorphism theorems” for AFP groups in Bass-Serre theory. Our main “subalgebra theorem” shows that, under rather general conditions, any von Neumann subalgebra $Q \subset M$ with the *relative property* (T) in the sense of ([P5]) (also called a *rigid inclusion*), can be conjugated by an inner automorphism of M into either M_1 or M_2 . We derive several “isomorphism theorems” in the case the amalgamation is over the scalars, $B = \mathbb{C}$, over a common Cartan subalgebra, $B = A$, or over a regular hyperfinite subfactor, $B = R$. The typical such statement shows that if $\theta : M \simeq N^t$ is an isomorphism from an AFP factor $M = M_1 *_B M_2 *_B \dots *_B M_n$ onto the amplification by some $t > 0$ of an AFP factor $N = N_1 *_C N_2 *_C \dots *_C N_n$, $1 \leq m, n \leq \infty$, with each M_i, N_j containing a “large” subalgebra with the relative property (T), then $m = n$ and $\theta(B \subset M_i)$ is unitarily conjugate to $(C \subset N_i)^t$, $\forall i$, after some permutation of indices.

When applied to the case $B = R$, these results allow us to obtain the first explicit calculations of outer automorphism groups of II_1 factors and answer in the affirmative a problem posed by A. Connes in 1973, on whether there exist II_1 factors M with no outer automorphism, i.e. with $\text{Out}(M) \stackrel{\text{def}}{=} \text{Aut}(M)/\text{Int}(M) = \{1\}$. More precisely, we show that if a group Γ is the free product of two infinite property (T) groups with no non-trivial characters, for example $\Gamma = SL(n_0, \mathbb{Z}) * SL(n_1, \mathbb{Z})$, $n_0, n_1 \geq 3$, then there exist actions of Γ on the hyperfinite II_1 factor R such that the corresponding crossed product factors $M = R \rtimes \Gamma$ have both trivial fundamental group, $\mathcal{F}(M) = \{1\}$, and trivial outer automorphism group, $\text{Out}(M) = \{1\}$. In fact, the general result shows that for any separable compact abelian group K there exist factors M with $\mathcal{F}(M) = \{1\}$ and $\text{Out}(M) = K$.

In turn, when applied to the case of amalgamated free products over a common Cartan subalgebra, our “isomorphism theorem” provides a Bass-Serre type result for *orbit equivalence* (OE) of actions of free product groups $\Gamma = \Gamma_1 * \dots * \Gamma_n$, $\Lambda = \Lambda_1 * \dots * \Lambda_m$ on the probability space. Thus, we show that if each Γ_i, Λ_j has an infinite normal subgroup with the relative property (T) of Kazhdan-Margulis (for instance Γ_i, Λ_j Kazhdan groups, $\forall i, j$) and $(\sigma, \Gamma), (\theta, \Lambda)$ are free, probability measure preserving (m.p.) actions with $\sigma|_{\Gamma_i}, \theta|_{\Lambda_j}$ ergodic $\forall i, j$, then $\sigma \sim_{OE} \theta$ implies $m = n$ and $\sigma|_{\Gamma_i} \sim_{OE} \theta|_{\Lambda_i}$, $\forall i$, after a permutation of the indices i . Note that the opposite implication holds true for arbitrary groups Γ_i, Λ_j , as shown by D. Gaboriau in ([G2]). In fact, we derive the component by component OE of actions under the weaker assumption that the group measure space factors associated with $(\sigma, \Gamma), (\theta, \Lambda)$ are stably isomorphic, i.e. when σ, θ are *von Neumann equivalent* (vNE). We use this vNE Bass-Serre rigidity and ([MS], [Ge1,2], [Fu1,2], [P6,8]) to give examples of group measure space factors M from free ergodic m.p. actions σ of free product groups $\Gamma = \Gamma_1 * \Gamma_2 * \dots$ such that $\mathcal{F}(M) = \{1\}$ and $\text{Out}(M) = H^1(\sigma, \Gamma)$, with explicit calculation of the abelian group $H^1(\sigma, \Gamma)$.

Finally, when applied to the case $B = \mathbb{C}$ our results become von Neumann algebra analogue of Kurosh’s classical theorems for free products of groups, similar to N. Ozawa’s

recent results of this type in ([O2]), but covering a different class of factors than ([O2]) and allowing amplifications. For instance, we show that if $N_i, 2 \leq i \leq n, M_j, 2 \leq j \leq m$, are property (T) II_1 factors in the sense of Connes-Jones (e.g. N_i, M_j group factors associated with Kazhdan groups, [CJ]) then $M_1 * M_2 * \dots * M_m \stackrel{\theta}{\simeq} (N_1 * N_2 * \dots * N_n)^t$ implies $m = n$ and $\theta(M_i)$ inner conjugate to $N_i^t, \forall i$, after some permutation of indices. In fact, in its most general form our result only requires M_i, N_j to be *weakly rigid* (*w-rigid*), i.e. to have diffuse, regular subalgebras with the relative property (T). Taking $M = N$ and $M_j = P^{s_j}$, with $\{s_j\}_j = S$ a multiplicative subgroup of $\mathbb{R}_+^*$ and P a w-rigid II_1 factor with trivial fundamental group (for instance the group factor $L(G)$ associated with $G = \mathbb{Z}^2 \rtimes SL(2, \mathbb{Z})$, cf. [P5]) and using a result of Dykema-Radulescu ([DyR]), we get $\mathcal{F}(M) = S$ for $M = *_{s \in S} P^s$. This provides a completely new class of factors with arbitrary given $S \subset \mathbb{R}_+^*$ as fundamental group than the ones in ([P6]). Indeed, the examples constructed in ([P6]) are group measure space factors, while the free group factors $*_{s \in S} P^s$ have no Cartan subalgebra, by results of Voiculescu ([V2]; see [Sh] and Remark 6.6 in this paper).

The key technical result behind all these applications is the above mentioned “subalgebra theorem”, of Bass-Serre type. We state it in details below, together with other main results in the paper, and also explain some of the ideas behind the proofs. An inclusion of finite von Neumann algebras $B \subset P$ will be called *homogeneous* if there exist $\{y_j\}_j \subset P$ with $E_B(y_i^* y_j) = \delta_{ij}, \forall i, j$ and $\Sigma_i y_j B$ dense in P . This technical assumption is satisfied by all inclusions coming from (cocycle) crossed products and (generalized) group measure constructions, or Cartan inclusions. It is also satisfied when P is an arbitrary finite von Neumann algebra and $B = \mathbb{C}$. Following ([P5]), a von Neumann subalgebra $Q \subset P$ has the *relative property* (T) (or $Q \subset P$ is a *rigid inclusion*) if any “deformation” of id_P by completely positive, sub-unital, sub-tracial maps, $\phi_n \rightarrow \text{id}_P$, is uniform on the unit ball of Q (see also [PeP]).

0.1. Theorem. *Let $(M_i, \tau_i), i = 1, 2$, be finite factors with a common von Neumann subalgebra $B \subset M_i$, such that $\tau_1|_B = \tau_2|_B$ and such that $B \subset M_i$ are homogeneous, $i = 1, 2$. Let $Q \subset M = M_1 *_B M_2$ be a diffuse von Neumann subalgebra with the relative property (T) such that no corner qQq of Q can be embedded into B . Then there exists a unique partition of 1 with projections q'_1, q'_2 in the commutant of Q in M such that $u_i(Qq'_i)u_i^* \subset M_i, i = 1, 2$, for some unitaries u_1, u_2 in M . Moreover, if the normalizer of Q in M generates a factor N , then there exists a unique $i \in \{1, 2\}$ such that $uQu^* \subset M_i$ for some $u \in \mathcal{U}(M)$, which also satisfies $uNu^* \subset M_i$.*

The proof of this result takes Sections 2 through 5 of the paper. It uses “deformation/rigidity” and “intertwining” techniques from ([P3,5,6]). Thus, we embed $M = M_1 *_B M_2$ into the larger algebra $\tilde{M} = M *_B (B \bar{\otimes} L(\mathbb{F}_2))$, whose abundance of deformations is used to show that “rigid parts” of M have to concentrate on certain subspaces with “bounded word-length”. This initial information is then used as a starting point in a word-reduction argument to obtain a Hilbert bimodule intertwining Q into one of the M_i ’s. The homogeneity condition is needed in order to measure the “size” of letters in M_i ’s. To get from this a unitary element conjugating Q into M_i , we prove in Section 1 a series of results on the

relative commutants and normalizers of subalgebras in AFP factors, using (2.1, 2.3 in [P6]).

If we take $B = \mathbb{C}$ in the Theorem 0.1 and use the fact that finite von Neumann algebras with Haagerup property have no diffuse subalgebras with the relative property (T), then we get an analogue of Kurosh's isomorphism theorem for free products of groups:

0.2. Theorem. *Let $(M_0, \tau_{M_0}), (N_0, \tau_{N_0})$ be finite von Neumann algebras with Haagerup's compact approximation property. Let $M_i, 1 \leq i \leq m$, and $N_j, 1 \leq j \leq n$, be w-rigid II_1 factors, where $m, n \geq 1$ are some cardinals (finite or infinite). If θ is an isomorphism of $M = *_{i=0}^m M_i$ onto N^t , where $N = *_{j=0}^n N_j$ and $t > 0$, then $m = n$ and, after some permutation of indices, $\theta(M_i)$ and N_i^t are unitary conjugate in N^t , $\forall i \geq 1$.*

Ozawa's pioneering result of this type in ([O2]) concerns free products of group factors $M_i = L(\Gamma_i), N_i = L(\Lambda_i)$ with each Γ_i, Λ_i a product of two or more ICC groups, either word hyperbolic (at least one of them) or amenable, typical examples being the groups $\mathbb{F}_{n_i} \times S_\infty$, not covered by 0.2 above. In turn, our typical M_i, N_i are factors from property (T) (more generally w-rigid) groups.

Letting $M_i = N_i, \forall i$, and $m < \infty$ in Theorem 0.2, it follows that if $\mathcal{F}(M_i) = \{1\}$ for some $1 \leq i \leq m$ (for example if $M_i = L(\mathbb{Z}^2 \rtimes \mathbb{F}_k)$, with $\infty > k \geq 2$, cf. [P5]), then $\mathcal{F}(M) = \{1\}$. Moreover, taking $m = \infty$ in 0.2 and using the “compression formula” for free products of infinitely many II_1 factors $(*_i M_i)^t \simeq *_i M_i^t$ in ([DyR]), we can include specific numbers into the fundamental group. Thus we get:

0.3. Corollary. *1°. Let $m \in \mathbb{N}$ and let $M_1, \dots, M_m$ be w-rigid II_1 factors. Let (M_0, τ_{M_0}) be a finite von Neuman algebra with Haagerup's compact approximation property. If one of the factors $M_i, m \geq i \geq 1$, has trivial fundamental group then so does $M = *_{i=0}^m M_i$.*

2°. If $S \subset \mathbb{R}_+^$ is an arbitrary infinite (possibly uncountable) subgroup and P is a w-rigid II_1 factor with trivial fundamental group (e.g. $P = L(\mathbb{Z}^2 \rtimes SL(2, \mathbb{Z}))$) then the II_1 factor $*_{s \in S} P^s$ has fundamental group equal to S .*

Since a group measure space factor $M = L^\infty(X, \mu) \rtimes_\sigma (\Gamma_1 * \Gamma_2)$ associated with a free ergodic m.p. action $(\sigma, \Gamma_1 * \Gamma_2)$ on a probability space (X, μ) can alternatively be viewed as an AFP factor $M = M_1 *_A M_2$, where $A = L^\infty(X, \mu)$, $M_i = A \rtimes_{\sigma|_{\Gamma_i}} \Gamma_i$, Theorem 0.1 allows us to obtain Bass-Serre type vNE and OE rigidity results for actions of free products of groups, as follows:

0.4. Theorem (vNE Bass-Serre rigidity). *Let Γ_0, Λ_0 be groups with Haagerup property and $\Gamma_i, \Lambda_j, 1 \leq i \leq n \leq \infty, 1 \leq j \leq m \leq \infty$, be ICC groups having normal, non virtually abelian subgroups with the relative property (T). Assume either Γ_0 is infinite or $n \geq 2$. Let σ (resp. θ) be a free ergodic m.p. action of $\Gamma = \Gamma_0 * \Gamma_1 * \dots$ (resp. $\Lambda = \Lambda_0 * \Lambda_1 * \dots$) on the probability space (X, μ) (resp. (Y, ν)) such that $\sigma_j = \sigma|_{\Gamma_j}$ (resp. $\theta_j = \theta|_{\Lambda_j}$) is ergodic $\forall j \geq 1$. Denote $M = L^\infty(X, \mu) \rtimes_\sigma \Gamma, N = L^\infty(Y, \nu) \rtimes_\theta \Lambda, M_i = L^\infty(X, \mu) \rtimes_{\sigma_i} \Gamma_i \subset M, N_j = L^\infty(Y, \nu) \rtimes_{\theta_j} \Lambda_j \subset N$ the corresponding group measure space factors. If $\alpha : M \simeq N^t$ is an isomorphism, for some $t > 0$, then $m = n$ and there exists a permutation π of indices $j \geq 1$ and unitaries $u_j \in N^t$ such that $\text{Ad}(u_j)(\alpha(M_j)) = N_{\pi(j)}^t, \text{Ad}(u_j)(\alpha(L^\infty(X, \mu))) = (L^\infty(Y, \nu))^t, \forall j \geq 1$. In particular, $\mathcal{R}_\sigma \simeq \mathcal{R}_\theta^t$ and $\mathcal{R}_{\sigma_j} \simeq \mathcal{R}_{\theta_{\pi(j)}}^t, \forall j \geq 1$.*

In particular, taking the isomorphism α between the group measure space factors in 0.4 to come from an orbit equivalence of the actions, one gets:

0.5. Corollary (OE Bass-Serre rigidity). *Let Γ_i, Λ_j , $0 \leq i \leq n \leq \infty$, $1 \leq j \leq m \leq \infty$, σ, θ be as in 0.4. If $\mathcal{R}_{\sigma, \Gamma} \simeq \mathcal{R}_{\theta, \Lambda}^t$ then $n = m$ and there exists a permutation π of the set of indices $i \geq 1$ such that $\mathcal{R}_{\sigma_i, \Gamma_i} \simeq \mathcal{R}_{\theta_{\pi(i)}, \Lambda_{\pi(i)}}^t$, $\forall i \geq 1$.*

Like in ([P6II]), the terminology “vNE rigidity” is used here in a broad sense, in the same spirit the terminology “OE rigidity” is being used in orbit equivalence ergodic theory ([Z1], [Fu1], [MS], [S]). It can designate results which from an isomorphism of group measure space factors derives orbit equivalence of the actions involved (“vNE/OE rigidity”, like 6.2 in [P5]), or even conjugacy of the actions (“vNE strong rigidity”, e.g. 7.1 in [P6II]). Theorem 0.4 brings out a new type of vNE rigidity, which we have labeled “Bass-Serre” because of its analogy to group theory results. It is a “vNE/OE”-type result but stronger, as it derives not only the orbit equivalence of the “main actions” (σ, Γ) , (θ, Λ) , but also the component by component orbit equivalence of their restrictions (σ_i, Γ_i) , (θ_i, Λ_i) .

The “vNE Bass-Serre rigidity” can be used in combination with OE rigidity results in orbit equivalence ergodic theory to get more insight on the group measure space factors involved. Thus, taking $\Gamma_0 = \Lambda_0 = \{1\}$ and $2 \leq n, m < \infty$ in 0.4, by Gaboriau’s results in ([G1]) it follows that the ℓ^2 -Betti numbers of Γ_i, Λ_j must satisfy $\beta_k(\Gamma_i) = \beta_k(\Lambda_i)/t$, $\forall 1 \leq i \leq n = m$, and $\sum_i \beta_k(\Lambda_i) + (n - 1) = (\sum_i \beta_k(\Gamma_i) + (n - 1))/t$, forcing $t = 1$. Also, if we take $\Gamma = \Gamma_0 * \Gamma_1 * \dots$ and σ as in 0.4 and add the conditions $\text{Out}(\mathcal{R}_{\sigma_1}) = \{1\}$ and (σ_1, Γ_1) not OE to (σ_i, Γ_i) , $\forall i \neq 1$, then $\text{Out}(\mathcal{R}_\sigma) = \{1\}$ and $\text{Out}(M) = H^1(\sigma, \Gamma)$. Examples of actions (σ_1, Γ_1) with the associated orbit equivalence relation $\mathcal{R}_{\sigma_1, \Gamma_1}$ having trivial outer automorphism group are constructed in ([Ge2], [Fu2], [MS]) and we construct some more, using the Monod-Shalom rigidity theorem ([MS]). The group $H^1(\sigma, \Gamma)$ can in turn be calculated by using ([P8]), thus getting explicit computations of $\text{Out}(M)$ for the group measure space factors M . The fact that one can choose the action (σ, Γ) to be free yet have restrictions $\sigma|_{\Gamma_i}$ isomorphic to specific Γ_i -actions, $\forall i$, is a consequence of ([To]) but we include a proof for the reader’s convenience (see 7.3 and A.1).

Theorem 0.1 is in fact used to obtain another (genuine) “vNE/OE rigidity” result in this paper, for free ergodic m.p. actions (σ, Γ) with Γ a free product of infinite groups, $\Gamma = \Gamma_0 * \Gamma_1$, and σ satisfying the relative property (T) of (5.10 in [P5]), i.e. such that $L^\infty(X, \mu) \subset L^\infty(X, \mu) \rtimes_\sigma \Gamma$ is a rigid inclusion. We recover this way the uniqueness of the HT Cartan subalgebra in the group-factors $L(\mathbb{Z}^2 \rtimes \mathbb{F}_n)$ and their amplifications, one of the main results in ([P5]).

Similarly, we obtain rigidity results for crossed product factors $M = R \rtimes_\sigma (\Gamma_0 * \Gamma_1)$ corresponding to actions $(\sigma, \Gamma_0 * \Gamma_1)$ on the hyperfinite II_1 factor R , by regarding M as an AFP factor $M = (R \rtimes \Gamma_0) *_R (R \rtimes \Gamma_1)$. In fact, in this case we can control even better the groups of symmetries $\mathcal{F}(M)$, $\text{Out}(M)$, with complete calculations. To state this result, let $f\mathcal{T}_R$ denote the class of actions $(\sigma, \Gamma_0 * \Gamma_1)$ of free product groups $\Gamma_0 * \Gamma_1$ on the hyperfinite factor R , satisfying the properties: (a). Γ_0 is free indecomposable; (b). Γ_1 is w-rigid (e.g. Γ_1 infinite Kazhdan group); (c). $R \subset R \rtimes_\sigma \Gamma_0$ is a rigid inclusion; (d). $\sigma|_{\Gamma_1}$ is a non-commutative

Bernoulli Γ_1 -action, i.e. R can be represented in the form $R = \overline{\otimes}_{g \in \Gamma_1} (M_{n \times n}(\mathbb{C}), tr)_g$, $n \geq 2$, with $\sigma|_{\Gamma_1}$ acting on it by left Bernoulli shifts; (e). $\sigma|_{\Gamma_1}$ is freely independent with respect to the normalizer $\mathcal{N}_0$ of $\sigma(\Gamma_0)$ in $\text{Out}(R)$.

To show that such actions exist, we first prove that for any two countable sets of automorphisms S_1, S_2 of R there exist $\theta \in \text{Aut}(R)$ such that S_1 and $\theta S_2 \theta^{-1}$ are “freely independent” (see 8.2 and A.2). Combining this with results from ([Ch], [P5], [NPS], [Va]) we deduce that for many arithmetic groups Γ_0 (in particular for $\Gamma_0 = SL(n, \mathbb{Z}), \forall n \geq 2$) and any w-rigid group Γ_1 , there exist actions $(\sigma, \Gamma_0 * \Gamma_1)$ on R in the class $f\mathcal{T}_R$. Using Theorem 0.1, properties (a) – (e) above and [P3], we get:

0.6. Theorem. *For any $\Gamma_0 = SL(n_0, \mathbb{Z}), n_0 \geq 2$ and any w-rigid group Γ_1 there exist actions σ of $\Gamma_0 * \Gamma_1$ on R in the class $f\mathcal{T}_R$. If $(\sigma, \Gamma_0 * \Gamma_1)$ is a $f\mathcal{T}_R$ action and we denote $M = R \rtimes_{\sigma} (\Gamma_0 * \Gamma_1)$, then $\mathcal{F}(M) = \{1\}$ and $\text{Out}(M) = \text{Char}(\Gamma_0) \times \text{Char}(\Gamma_1)$.*

0.7. Corollary. *Given any compact abelian group K , there exist separable II_1 factors M with $\mathcal{F}(M) = \{1\}$ and $\text{Out}(M) = K$. For instance, if $(\sigma, \Gamma_0 * \Gamma_1)$ is a $f\mathcal{T}_R$ action and $M = R \rtimes_{\sigma} (\Gamma_0 * \Gamma_1)$ is the associated crossed product factor, with $\Gamma_0 = SL(n, \mathbb{Z})$, $\Gamma_1 = SL(m, \mathbb{Z}) \times \hat{K}$ for some $n, m \geq 3$, then $\mathcal{F}(M) = \{1\}$ and $\text{Out}(M) = K$. Moreover, denoting $M^{\infty} = M \overline{\otimes} (\mathcal{B}(\ell^2 \mathbb{N}))$ the associated II_{∞} factor, we have $\text{Out}(M^{\infty}) = K$.*

The study of outer automorphisms of type II von Neumann factors was at the core of Connes decomposition theory for factors of type III and his classification of amenable factors, in the early 70’s ([C1,2]). Two subsequent seminal papers ([C3,4]) gave the first indications that the outer symmetry groups $\text{Out}(M)$ and $\mathcal{F}(M)$ can reflect rigidity properties of non-amenable factors. In particular, it was shown in ([C3]) that $\mathcal{F}(M)$ and $\text{Out}(M)$ are countable for group factors associated with ICC groups with the property (T). The recent rigidity results in ([P5,6,9]) provide explicit calculations of $\mathcal{F}(M)$ for large families of group measure space factors M , and reduce the calculation of $\text{Out}(M)$ to the computation of the commutants of the corresponding group actions. However, such commutants are difficult to compute, being left as an open problem even in the case of Bernoulli actions (see [P6II]). The calculation of $\text{Out}(M)$ that we obtain in this paper for crossed product factors arising from actions of free products of w-rigid groups on the probability space and on the hyperfinite factor thus give the first such explicit computations.

This work initiated while the authors were visiting the Laboratoire d’Algèbres d’Opérateurs at the Institut de Mathématiques of the University Paris 7 during the Fall of 2004. It is a pleasure for us to thank the CNRS and the members of the Lab for their support and kind hospitality. We are very grateful to Damien Gaboriau for illuminating discussions and comments on our work and Bass-Serre theory. We thank Dima Shlyakhtenko for kindly pointing out to us Remark 6.6.

1. CONJUGATING SUBALGEBRAS IN AFP FACTORS

1.1. AFP algebras. Let $(M_1, \tau_1), (M_2, \tau_2)$ be finite von Neumann algebras with a common von Neumann subalgebra $B \subset M_i, i = 1, 2$, such that $\tau_1|_B = \tau_2|_B$. We denote by

$(M_1 *_B M_2, \tau_1 * \tau_2)$ the finite von Neumann algebra *free product with amalgamation* (AFP) of $(M_1, \tau_1; B)$ and $(M_2, \tau_2; B)$, as defined in ([V1] and pp. 384-385 of [P4]). Thus, $M_1 *_B M_2$ has a dense $*$ -subalgebra

$$(1.1.1) \quad B \oplus \bigoplus_{n \geq 1} \bigoplus_{\substack{i_j \in \{1,2\} \\ i_1 \neq i_2, \dots, i_{n-1} \neq i_n}} \text{sp}(M_{i_1} \ominus B)(M_{i_2} \ominus B) \cdots (M_{i_n} \ominus B)$$

with the trace $\tau = \tau_1 * \tau_2$ defined on *reduced words* by $\tau(x) = \tau_1(x) = \tau_2(x)$ for $x \in B$ and $\tau(x) = 0$ for $x = x_{i_1} x_{i_2} \cdots x_{i_n}$, with $x_{i_k} \in M_{i_k} \ominus B$, $i_k \in \{1, 2\}$, $i_1 \neq i_2, \dots, i_{n-1} \neq i_n$. Thus, the vector subspaces B and $\text{sp}(M_{i_1} \ominus B)(M_{i_2} \ominus B) \cdots (M_{i_n} \ominus B) \subset M$ in the above sum are all mutually orthogonal with respect to the scalar product given by the trace τ . Also, their closure in $L^2(M, \tau)$ gives mutually orthogonal Hilbert B -bimodules,

$$L^2((M_{i_1} \ominus B)(M_{i_2} \ominus B) \cdots (M_{i_n} \ominus B)) \simeq \mathcal{H}_{i_1}^0 \otimes_B \mathcal{H}_{i_2}^0 \otimes_B \cdots \otimes_B \mathcal{H}_{i_n}^0,$$

summing up to $L^2(M, \tau)$, where $\mathcal{H}_i^0 = L^2(M_i) \ominus L^2(B)$.

1.2. Controlling intertwiners and relative commutants. In this sub-section we prove a very useful “dichotomy-type” result for subalgebras Q of AFP factors $M = M_1 *_B M_2$. It shows that if Q sits in one of the factors, say M_1 , then it can either be conjugated into the “core” B of the AFP algebra M , or else all its normalizer lies in M_1 , and even all “intertwining” Hilbert $Q - M_1$ bimodules $\mathcal{H} \subset L^2(M)$ with $\dim(\mathcal{H}_{M_1}) < \infty$ must be entirely contained in $L^2(M_1)$! Also, in this second alternative, any bimodule intertwining Q into the other factor, M_2 , vanishes.

The first results of this type have been obtained in ([P1]), in the case of “plain” free product factors, $M = M_1 * M_2$. The next theorem provides a sharp generalization of those results. The proof uses the basic “intertwining criteria” (2.1 and 2.3 in [P6]), following arguments similar to (3.1 in [P6]).

1.2.1. Theorem. *Let (M_1, τ_1) and (M_2, τ_2) be finite von Neumann algebras and B a common von Neumann subalgebra such that $\tau_1|_B = \tau_2|_B$. Let $M = M_1 *_B M_2$, $0 \neq q \in \mathcal{P}(M_1)$ and $Q \subset qM_1q$ a von Neumann subalgebra. Assume no corner of Q can be embedded into B inside M_1 , i.e. $Q' \cap q\langle M_1, B \rangle q$ contains no non-zero finite projections. If $0 \neq \xi \in L^2(qM)$ satisfies $Q\xi \subset L^2(\sum_i \xi_i M_k)$ for some $k \in \{1, 2\}$ and some $\xi_1, \dots, \xi_n \in L^2(M)$, then $k = 1$ and $\xi \in L^2(M_1)$. In particular, $Q' \cap qMq \subset M_1$, the normalizer $\mathcal{N}_{qMq}(Q)$ of Q in qMq is contained in qM_1q and if $x \in M$ satisfies $Qx \subset xM_2$ then $x = 0$.*

Proof. Let p denote the orthogonal projection of $L^2(M)$ onto the Hilbert subspace $\overline{Q\xi M_k}^{\|\cdot\|_2} \subset L^2(M)$. Note that $p \in Q' \cap q\langle M, e_{M_k} \rangle q$ and $0 \neq \text{Tr}(p) < \infty$, where $\text{Tr} = \text{Tr}_{\langle M, e_{M_k} \rangle}$ denotes the canonical trace on $\langle M, e_{M_k} \rangle$. To prove that $k = 1$ and $\xi \in M_k$ it is sufficient to show that $p \leq e_{M_k}$, equivalently $(1 - e_{M_k})p(1 - e_{M_k}) = 0$. Indeed, because then $Q\xi M_k \subset L^2(M_k)$, so in particular $\xi \in L^2(M_k)$ and $u\xi \in L^2(M_k)$, $\forall u \in \mathcal{U}(Q)$. But since no corner of Q can be embedded into B inside M_1 , by (2.3 in [P6]) it follows that for any $\varepsilon > 0$ there

exists $u \in \mathcal{U}(Q)$ such that $\|E_B(u)\|_2 \leq \varepsilon$. Thus, if $k = 2$ then $\xi, u\xi \in L^2(M_2)$ so that $u\xi = E_{M_2}(u\xi) = E_B(u)\xi$ and by the Cauchy-Schwartz inequality we have

$$\|\xi\|_1 = \|u\xi\|_1 = \|E_B(u)\xi\|_1 \leq \|E_B(u)\|_2 \|\xi\|_2 \leq \varepsilon \|\xi\|_2.$$

Since $\varepsilon > 0$ is arbitrary, this shows $\xi = 0$. Thus, the only possibility is that $k = 1$, i.e. $\xi \in L^2(M_1)$.

By taking spectral projections, to show that $(1 - e_{M_k})p(1 - e_{M_k}) = 0$ it is in fact sufficient to show that if $f \in Q' \cap \langle M, e_{M_k} \rangle$ is a projection such that $0 \neq \text{Tr}(f) < \infty$ and $f \leq 1 - e_{M_k}$, then $f = 0$. To this end, we'll show that $\|f\|_{2,Tr}$ is arbitrarily small.

Thus, let $\eta_0 = 1, \eta_1, \dots, \eta_n, \dots \subset M$ be an orthonormal basis of M over M_k , i.e. $E_{M_k}(\eta_i^* \eta_j) = \delta_{ij} p_j \in \mathcal{P}(M_k)$ and $\|\eta\| < \infty, \forall i, j$. If we denote $f_n = \sum_{i=1}^n \eta_i e_{M_k} \eta_i^*$ then, as f has finite trace and $f \leq 1 - e_{M_k} = \sum_{i=1}^\infty \eta_i e_{M_k} \eta_i^*$, there exists $n \in \mathbb{N}$ such that $\|f_n f - f\|_{2,Tr} < \epsilon \|f\|_{2,Tr}$. Thus, if $u \in \mathcal{U}(Q)$ then

$$(1.2.1) \quad \begin{aligned} \text{Tr}(f_n u f_n u^*) &\geq \text{Tr}(f f_n u f_n u^*) \\ &\quad - |\text{Tr}(f f_n (1 - f) u f_n u^*)| - |\text{Tr}((1 - f) f_n u f_n u^*)|. \end{aligned}$$

Using that $f_n f$ is ϵ -close to f in the norm $\|\cdot\|_{2,Tr}$ and that f commutes with $u \in Q$ we deduce:

$$(1.2.2) \quad \text{Tr}(f f_n u f_n u^*) = \text{Tr}(f_n u f_n u^*) \geq (1 - 2\epsilon - \epsilon^2) \|f\|_{2,Tr}^2.$$

Similarly we have:

$$(1.2.3) \quad |\text{Tr}(f f_n (1 - f) u f_n u^*)| + |\text{Tr}((1 - f) f_n u f_n u^*)| \leq 2\epsilon(1 + \epsilon) \|f\|_{2,Tr}^2.$$

Combining (1.2.1) – (1.2.3) we get

$$(1.2.4) \quad \text{Tr}(f_n u f_n u^*) \geq (1 - 4\epsilon - 3\epsilon^2) \|f\|_{2,Tr}^2, \forall u \in \mathcal{U}(Q).$$

On the other hand,

$$(1.2.5) \quad \text{Tr}(f_n u f_n u^*) = \text{Tr}(\sum_{i,j} \eta_i e_{M_k} \eta_i^* u \eta_j e_{M_k} \eta_j^* u^*) = \sum_{i,j} \|E_{M_k}(\eta_i u \eta_j^*)\|_2^2.$$

Thus, in order to prove that $\|f\|_{2,Tr}$ is small, it is sufficient to prove that $\forall \eta_0, \dots, \eta_n \in M \ominus M_k, \forall \epsilon > 0, \exists u \in \mathcal{U}(Q)$ such that $\|E_{M_k}(\eta_i u \eta_j^*)\|_2 \leq \epsilon, \forall 0 \leq i, j \leq n$. Furthermore, by (1.1.1) and Kaplansky's density theorem it is enough to prove this in the case η_i are reduced words of the form $\eta_i = \delta_i x_i$ such that one of the following holds true: (a). δ_i is a reduced word that ends with a letter in $M_2 \ominus B$ and x_i is either equal to 1 or contained in $M_1 \ominus B$; (b). $k = 2, \delta_i = 1$ and $x_i \in M_1 \ominus B$. Since $x_i u x_j^* \in M_1$, if we denote $y = x_i u x_j^* - E_B(x_i u x_j^*) \in M_1 \ominus B$ then in both cases (a) and (b) the reduced word $\delta_i y \delta_j^*$ is

perpendicular to M_k . Indeed, in case (a), $\delta_i y \delta_j^*$ lies in $\dots(M_{k'} \ominus B)(M_k \ominus B)(M_{k'} \ominus B)\dots$, where $\{k, k'\} = \{1, 2\}$, and thus it has length at least 3, so $\delta_i y \delta_j^* \perp M_k$ by (1.1.1). In case (b), $\delta_i y \delta_j^* \in M_1 \ominus B$ so it is perpendicular to $M_2 = M_k$. Using $\delta_i y \delta_j^* \perp M_k$ we then get:

$$\begin{aligned} E_{M_k}(\eta_i u \eta_j^*) &= E_{M_k}(\eta_i y \eta_j^*) + E_{M_k}(\delta_i E_B(x_i u x_j^*) \delta_j^*) \\ &= E_{M_k}(\delta_i E_B(x_i u x_j^*) \delta_j^*), \end{aligned}$$

implying that

$$\begin{aligned} \|E_{M_k}(\eta_i u \eta_j^*)\|_2 &= \|E_{M_k}(\delta_i E_B(x_i u x_j^*) \delta_j^*)\|_2 \\ &\leq \|\delta_i\| \|\delta_j\| \|E_B(x_i u x_j^*)\|_2. \end{aligned}$$

But by the hypothesis and (2.3 in [P6]), for any $\epsilon > 0$ we can find $u \in \mathcal{U}(Q)$ such that $\|E_B(x_i u x_j^*)\|_2 \leq \epsilon / (\|\delta_i\| \|\delta_j\|)$. $\square$

Note that under the conditions of the above theorem, not only the normalizer $\mathcal{N}_{qMq}(Q)$ of Q in qMq follows contained in M_1 , but also the normalizer of the von Neumann algebra generated by $\mathcal{N}_{qMq}(Q)$, and so on. In fact, even the unitaries $u \in qMq$ with the property that $uQu^* \cap qM_1q$ is not embeddable into B follow contained in M_1 . More generally, if $Q \subset qM_1q$ is a von Neumann subalgebra such that $Q' \cap q\langle M_1, B \rangle q$ contains no non-zero finite projections, and if we denote by $N_1 = N(Q, M_1; B)$ the von Neumann subalgebra of qMq generated by unitaries $u \in qMq$ such that $(uQu^* \cap M_1)' \cap q\langle M_1, B \rangle q$ contains no non-zero finite projections then $N_1 \subset M_1$. If we then repeat this operation, taking $N_2 = N(N_1, M_1; B)$ be the von Neumann algebra generated by all unitaries $u \in qMq$ such that $uN_1u^* \cap q\langle M_1, B \rangle q$ contains no non-zero finite projections, then $N_2 \subset M_1$. We can of course continue this procedure inductively until it “stops”, i.e. until we reach an N_i such that $N(N_i, M_1; B) = N_i$. More, formally, consider the following:

1.2.2. Definition. If $q \in \mathcal{P}(B)$ and $Q \subset qMq$ we consider by (transfinite) induction the strictly increasing family of von Neumann algebras $Q = N_0 \subset N_1 \subset \dots \subset N_j \subset \dots \subset N_i$, indexed by the first ι ordinals, such that: (a). For each $j < \iota$, $N_{j+1} = N(N_j, M_1; B)$ and $N_j \neq N_{j+1}$; (b). $N(N_i, M_1; B) = N_i$; (c). If $j \leq \iota$ has no “predecessor” then $N_j = \cup_{n < j} N_n$. We then denote $\tilde{N}(Q, M_1; B) = N_i$ and call it the *weak quasi normalizer* (wq-normalizer) of Q in qMq relative to $(M_1; B)$. Note that in fact both the definition of $N(Q, M_1; B)$ and $\tilde{N}(Q, M_1; B)$ make actually sense for any finite von Neumann algebra (M, τ) and von Neumann subalgebras $B, M_1 \subset M$, $Q \subset qMq$, with $q \in \mathcal{P}(M_1)$.

This definition is analogous to the definition of wq-normalizer of a subgroup $H \subset G$ used in ([P5,6,8]). It is easy to see that $\tilde{N}(Q, M_1; B)$ is the smallest von Neumann subalgebra P of qMq such that $(uPu^* \cap qM_1q)' \cap q\langle M, e_B \rangle q$ contains non-zero finite projections $\forall u \in (qMq) \setminus P$. Theorem 1.1 thus implies:

1.2.3. Corollary. *With (M_i, τ_i) , $q \in B \subset M_i$, $i = 1, 2$, $M = M_1 *_B M_2$ as in 1.2.1, let $Q \subset qM_1q$ be a von Neumann subalgebra such that no corner of Q can be embedded into B inside M_1 , i.e. $Q' \cap q\langle M_1, B \rangle q$ contains no non-zero finite projections. Then $\tilde{N}(Q, M_1; B) \subset M_1$.*

We will make repeated use of the following application of 1.2.1, which shows that if one of the algebras M_i involved in an amalgamated free product $M = M_1 *_B M_2$ contains a regular subalgebra Q , then Q must necessarily be contained in B , modulo inner conjugacy:

1.2.4. Corollary. *Let (M_i, τ_i) , $B \subset M_i$, $i = 1, 2$, be as in 1.1 and denote $M = M_1 *_B M_2$. Let $Q \subset qM_1q$ be a von Neumann subalgebra, for some $q \in \mathcal{P}(B)$ with $qBq \neq qM_2q$. Assume Q is regular in qMq . Then one can embed a corner of Q into B inside M_1 , i.e. $Q' \cap q\langle M_1, e_B \rangle q$ contains non-zero finite-trace projections.*

Proof. If $Q' \cap q\langle M_1, e_B \rangle q$ contains no non-zero finite-trace projection then by Theorem 1.1 the normalizer $\mathcal{N}_M(Q)$ of Q in qMq is contained in M_1 . Since $\mathcal{N}_M(Q)'' = qMq$, this implies $qMq = qM_1q$, thus $qM_2q = qBq$, a contradiction. $\square$

1.3. Locating subalgebras by means of normalizers. In this and the next sub-section we prove that if a subalgebra of $M = M_1 *_B M_2$ is normalized by “many” unitaries in M_1 then it must necessarily be contained in M_1 . This technical result will in fact not be needed until Section 7, where it plays a key role in the proof of the Bass-Serre type Theorem 7.4. The proof uses the intertwining criteria in ([P6]) and a careful asymptotic analysis of elements written in AFP expansion (1.1.1). We first prove the result assuming the subalgebra we want to “locate” is a unitary conjugate of a subalgebra of B . This assumption will be shown to be redundant in 1.4, in the case $B = A$ is Cartan in M .

1.3.1. Proposition. *Let Λ_1, Λ_2 be discrete groups and $\sigma : \Lambda \rightarrow \text{Aut}(B, \tau)$ an action of $\Lambda = \Lambda_1 * \Lambda_2$ on the finite von Neumann algebra (B, τ) . Denote $M = B \rtimes_\sigma \Lambda = M_1 *_B M_2$, where $M_i = B \rtimes_{\sigma|_{\Lambda_i}} \Lambda_i$, $i = 1, 2$. Let $q \in \mathcal{P}(B)$, $B_0 \subset qBq$ a von Neumann subalgebra, $u \in \mathcal{U}(qMq)$ and denote $\mathcal{N} = \{v \in \mathcal{U}(qM_1q) \mid v(uqB_0qu^*)v^* = uqB_0qu^*\}$. Assume no corner of $\mathcal{N}''$ can be embedded into B inside M_1 . Then $uqB_0qu^* \subset M_1$.*

Proof. Assume there exists $b_0 \in qB_0q$, $\|b_0\| \leq 1$ such that $d_0 = ub_0u^* - E_{M_1}(ub_0u^*)$ satisfies $c = \|d_0\|_2 > 0$. To get a contradiction, we first show that all the unit ball of $uqBqu^*$ can be embedded into a set of the form $\Sigma_{g \in F}(B)_1u_g$ with $F \subset \Lambda$ finite, $(B)_1$ denoting the unit ball of B . We need the following:

1.3.2. Lemma. *Let (B, τ) be a finite von Neumann algebra, $\sigma : \Lambda \rightarrow \text{Aut}(B, \tau)$ an action, $M = B \rtimes_\sigma G$ the corresponding crossed product finite von Neumann algebra and $\{u_g\}_g \subset M$ the canonical unitaries. For any finite set in the unit ball of M , $S_0 \subset (M)_1$, and any $\varepsilon > 0$, there exists $F \subset \Lambda$ finite such that $x(B)_1y^* \subset_\varepsilon \Sigma_{g \in F}(B)_1u_g$, $\forall x, y \in S_0$.*

Proof. By Kaplansky’s density theorem there exists a finite set $F_0 \subset \Lambda$ and elements $\{b_g^x \in B \mid x \in S_0, g \in F_0\}$, such that $x_0 = \Sigma_{g \in F_0} b_g^x u_g$ satisfies $\|x_0\| \leq 1$ and $\|x - x_0\|_2 \leq \varepsilon/2$, $\forall x \in S_0$. If we put $F = F_0 F_0^{-1}$ then we clearly have $x_0 B y_0^* \subset \Sigma_{g \in F} B u_g$, $\forall x, y \in S_0$. On the other hand, if $b \in B$ satisfies $\|b\| \leq 1$ and we let $x_0 b y_0^* = \Sigma_g b_g u_g$ then $b_g = E_B((x_0 b y_0^*) u_g^*)$ and thus $\|b_g\| \leq \|(x_0 b y_0^*) u_g^*\| \leq 1$. This implies $\|x b y^* - x_0 b y_0^*\|_2 \leq \varepsilon$ and thus $x b y^* \subset_\varepsilon \Sigma_{g \in F}(B)_1 u_g$. $\square$

By Lemma 1.3.2, it follows that there exists $F \subset \Lambda$ finite such that $uq(B)_1qu^* \subset_{\varepsilon/2} \Sigma_{g \in F}(B)_1u_g$, where $\varepsilon = c^2/4$. Denote $N = \mathcal{N}'' \subset qM_1q$.

For any $v \in \mathcal{N} \subset qM_1q$ we then have $v(ub_0u^*)v^* \in_{\varepsilon/2} \Sigma_{g \in F}(B)_1u_g$ as well. Since $v(ub_0u^*)v^* = vd_0v^* + v(E_{M_1}(ub_0u^*))v^*$, with $vd_0v^* \perp M_1$ and $v(E_{M_1}(ub_0u^*))v^* \in M_1$, by Pythagora it follows that $vd_0v^* \in_{\varepsilon/2} \Sigma_{g \in F_0}(B)_1u_g$, where $F_0 = F \setminus \Lambda_1$. Letting $d_1 \in \Sigma_{g \in F_0}(B)_1u_g$ be so that $\|d_0 - d_1\|_2 \leq \varepsilon/2$, we have thus shown:

(1.3.3). $\exists F_0 \subset \Lambda \setminus \Lambda_1$ finite and $d_1 \in \Sigma_{g \in F_0}(B)_1u_g$ with $\|d_1\| \leq |F_0|$, $\|d_1\|_2 \geq c - c^2/8 > 0$, such that $vd_1v^* \in_{\varepsilon} \Sigma_{g \in F_0}(B)_1u_g$, $\forall v \in \mathcal{N}$, where $\varepsilon = c^2/4$.

Now note that by the condition satisfied by the algebra $\mathcal{N}'' = N$, from (2.3 in [P6]) it follows that:

(1.3.4) $\forall K \subset \Lambda_1$ finite and $\delta > 0$, $\exists v \in \mathcal{N}$ such that if ξ denotes the projection of v onto the Hilbert space $\oplus_{h \in \Lambda_1 \setminus K} L^2(B)u_h$ then $\|v - \xi\|_2 \leq \delta$.

At this point, we need the following:

1.3.5. Lemma. *With (B, τ) , (σ, Λ) , $M, \{u_g\}_g$ as in 1.3.2, assume $\Lambda = \Lambda_1 * \Lambda_2$. Let $F_0 \subset \Lambda \setminus \Lambda_1$ be a finite set. There exists $K = K(F_0) \subset \Lambda_1$ finite such that any $\xi \in L^2(B \rtimes \Lambda_1)$ supported by $\Lambda_1 \setminus K$ satisfies $\xi(\Sigma_{g \in F_0}Bu_g)M_1 \perp \Sigma_{g \in F_0}Bu_g$.*

Proof. Note that each irreducible alternating word $g \in F_0$ has at least one letter from Λ_2 . Let K_0 denote the set of elements in Λ_1 that can appear as first letter in a word g in F_0 (including the trivial letter e) and denote $K = K_0K_0^{-1}$. Then $(\Lambda_1 \setminus K)K_0 \cap K_0 = \emptyset$. Now note that if $\xi \in L^2(B \rtimes \Lambda_1)$ is supported by $\Lambda_1 \setminus K$ then any element η in $\xi(\Sigma_{g \in F}Bu_g)$ is supported on elements $g \in G$ that begin by a letter in $(\Lambda_1 \setminus K)K_0 \subset \Lambda_1 \setminus K_0$. Moreover, this is still the case for elements of the form ηx for $x \in M_1$. In turn, any g in the support of an element in $(\Sigma_{g \in F_0}Bu_g)$ begins by a letter in K_0 . Thus, the two vector spaces $\xi(\Sigma_{g \in F_0}Bu_g)M_1$, $(\Sigma_{g \in F_0}Bu_g)$ are supported on disjoint subsets of $\Lambda_1 * \Lambda_2$ and are thus perpendicular. $\square$

We continue the proof of Proposition 1.3.1 by letting $K = K(F_0)$ be given by 1.3.5, for the finite set $F_0 \subset \Lambda \setminus \Lambda_1$ from (1.3.3). Let $\delta = \varepsilon/|F_0|$ and choose $\xi \in L^2(B \rtimes \Lambda_1)$ supported on $\Lambda_1 \setminus K$, as given by (1.3.4). Then

$$\|\xi d_1v^* - vd_1v^*\|_2 \leq \|\xi - v\|_2\|d_1\| \leq \delta|F_0| \leq \varepsilon,$$

which together with (1.3.3) implies $\xi d_1v^* \in_{2\varepsilon} \Sigma_{g \in F_0}Bu_g$. But by Lemma 1.3.5 we have $\xi d_1v^* \perp \Sigma_{g \in F_0}Bu_g$. Thus $\|\xi d_1v^*\|_2 \leq 2\varepsilon$. On the other hand

$$\|\xi d_1v^*\|_2 = \|\xi d_1\|_2 \geq \|d_1\|_2 - \|\xi - v\|_2\|d_1\| \geq c - c^2/8 - \varepsilon > 2\varepsilon,$$

a contradiction which ends the proof of Proposition 1.3.1. $\square$

1.4. A Cartan conjugacy result. We now prove that if the “core” B of an AFP algebra $M = M_1 *_B M_2$ is maximal abelian and regular (thus Cartan) in M , then any other Cartan subalgebra $A_0 \subset M$ which is normalized by “many” unitaries in M_1 is a unitary conjugate of $B = A$. Note that it strenghtens both Corollary 1.2.4 and Proposition 1.3.1, in the case the core $B = A$ is abelian and Cartan in M .

1.4.1 Theorem. *Let Λ_1, Λ_2 be infinite discrete groups and $\sigma : \Lambda \rightarrow \text{Aut}(A, \tau)$ a free ergodic action of $\Lambda = \Lambda_1 * \Lambda_2$ on a diffuse abelian von Neumann algebra (A, τ) . Denote $M = A \rtimes_{\sigma} \Lambda = M_1 *_A M_2$, where $M_i = A \rtimes_{\sigma|_{\Lambda_i}} \Lambda_i$, $i = 1, 2$. Let $q \in \mathcal{P}(A)$, $A_0 \subset qMq$ a Cartan subalgebra such that no corner of $(\mathcal{N}_{qMq}(A_0) \cap qM_1q)''$ can be embedded into A inside M_1 . Then $A_0 \subset qM_1q$ and there exists $u \in \mathcal{U}(qM_1q)$ such that $uA_0u^* = Aq$.*

1.4.2. Lemma. *Let Λ_1, Λ_2 be discrete groups and let $\sigma : \Lambda = \Lambda_1 * \Lambda_2 \rightarrow \text{Aut}(B, \tau)$ be a trace preserving action on a finite von Neumann algebra (B, τ) . Denote $M = B \rtimes_{\sigma} \Lambda = M_1 *_B M_2$ and let $M_1 = B \rtimes_{\sigma|_{\Lambda_1}} \Lambda_1 \subset M$. Let $q \in \mathcal{P}(B)$ and assume that $A_0 \subset qMq$ is a diffuse abelian von Neumann subalgebra such that no corner of A_0 can be embedded into qM_1q inside M . Then for any $\varepsilon > 0$ there exist $F \subset \Lambda \setminus \Lambda_1$ finite and $x_1, x_2 \in \Sigma_{g \in F} Bu_g$ such that any $u \in \mathcal{N}_{qMq}(A_0)$ satisfies*

$$\|ux_1u^*x_2 - x_2ux_1u^*\|_2 \leq \varepsilon, \sqrt{\tau(q)} - \varepsilon \leq \|ux_1u^*x_2\|_2 \leq \sqrt{\tau(q)} + \varepsilon.$$

Proof. By the assumption on A_0 it follows from (2.3 in [Po6]) that there exists $a_1 \in \mathcal{U}(A_0)$ such that $\|E_{M_1}(a_1)\|_2 < \varepsilon/4$. Thus, we can find $F_1 \subset \Lambda \setminus \Lambda_1$ finite and $x_1 \in \Sigma_{g \in F_1} Bu_g$ such that $\|a_1 - x_1\|_2 \leq \varepsilon/4$. Repeating the above argument we can now find $a_2 \in \mathcal{U}(A_0)$, a finite set F with $F_1 \subset F \subset \Lambda \setminus \Lambda_1$ and $x_2 \in \Sigma_{g \in F} Bu_g$ such that $\|a_2 - x_2\|_2 \leq \varepsilon/(4\|x_1\|)$. Using these inequalities we get for $u \in \mathcal{N}_{qMq}(A_0)$ (in fact for all $u \in M$ with $\|u\| \leq 1$):

$$\begin{aligned} \|ux_1u^*x_2 - ua_1u^*a_2\|_2 &\leq \|ux_1u^*(x_2 - a_2)\|_2 + \|u(x_1 - a_1)u^*a_2\|_2 \\ &\leq \|x_1\| \|x_2 - a_2\|_2 + \|x_1 - a_1\|_2 \leq \|x_1\| \varepsilon/(4\|x_1\|) + \varepsilon/4 = \varepsilon/2, \forall u \in (M)_1. \end{aligned}$$

Similarly, it follows that $\|x_2ux_1u^* - a_2ua_1u^*\|_2 \leq \varepsilon/2, \forall u \in (M)_1$. Finally, if $u \in \mathcal{N}_{qMq}(A_0)$ then $ua_1u^*a_2 = a_2ua_1u^*$ (because A_0 is abelian) and $\|ua_1u^*a_2\|_2 = \sqrt{\tau(q)}$, and thus by combining this with the above inequalities we get the desired estimates. $\square$

1.4.3. Lemma. *With the same notations and assumptions as in 1.4.2, let $F \subset \Lambda \setminus \Lambda_1$ be a finite set and let $x_1, x_2 \in \Sigma_{g \in F} Bu_g$. Then $\forall \varepsilon > 0, \exists K = K(F, \varepsilon) \subset \Lambda_1$ finite and $\delta = \delta(F, \varepsilon) > 0$ such that if $u \in (M_1)_1$ satisfies $\|E_B(uu_g^*)\|_2 \leq \delta, \forall g \in K$, then it must also satisfy $ux_1u^*x_2 \perp_{\varepsilon} x_2ux_1u^*$.*

Proof. Since $F \subset \Lambda \setminus \Lambda_1$ is finite, by the free decomposition $\Lambda = \Lambda_1 * \Lambda_2$ one readily deduces that there exists $K = K^{-1} \subset \Lambda_1$ finite such that $(\Lambda_1 \setminus K)F(\Lambda_1 \setminus K)F$ has empty intersection with $F(\Lambda_1 \setminus K)F(\Lambda_1 \setminus K)$. Next, let $u \in (M_1)_1$ and set $u' = \Sigma_{g \in K} E_B(uu_g^*)u_g$ and $u'' = u - u'$. Then u'' is supported on $\Lambda_1 \setminus K$ and we have the decomposition

$$ux_1u^*x_2 = u''x_1u''^*x_2 + u'x_1u^*x_2 + ux_1u'^*x_2 - u'x_1u'^*x_2.$$

Denote $x_{12} = u''x_1u''^*x_2$. Then, x_{12} is supported on $(\Lambda_1 \setminus K)F(\Lambda_1 \setminus K)F$ and we have the following estimate:

$$\|ux_1u^*x_2 - x_{12}\|_2 \leq (2\|u\| + \|u'\|)\|x_1\|\|x_2\|\|u'\|_2$$

$$\leq (2 + |K|)\|x_1\|\|x_2\|\|u'\|_2$$

Similarly, if we denote $x_{21} = x_2 u'' x_1 u''^*$, then x_{21} is supported on $F(\Lambda_1 \setminus K)F(\Lambda_1 \setminus K)$ and $\|x_2 u x_1 u^* - x_{21}\|_2 \leq (2 + |K|)\|x_1\|\|x_2\|\|u'\|_2$.

Next, we show that K and $\delta = \varepsilon(12|K|(\|x_1\|\|x_2\| + 1))^{-3/2}$ satisfy the conclusion. To this end, let $u \in (M_1)_1$ be such that $\|E_B(uu_g^*)\|_2 \leq \delta, \forall g \in K$. Then

$$\|u'\|_2 = (\sum_{g \in K} \|E_B(uu_g^*)\|_2^2)^{1/2} \leq \varepsilon|K|^{-1}(12(\|x_1\|\|x_2\| + 1))^{-3/2},$$

hence $\|ux_1 u^* x_2 - x_{12}\|_2 \leq \varepsilon(\|x_1\|\|x_2\| + 1)^{-1}/4$ and $\|x_2 u x_1 u^* - x_{21}\|_2 \leq \varepsilon(\|x_1\|(\|x_2\| + 1))^{-1}/4$. Also, we have:

$$\|x_{12}\|_2 \leq \|x_{12} - ux_1 u^* x_2\|_2 + \|ux_1 u^* x_2\|_2$$

$$\leq \varepsilon(\|x_1\|\|x_2\| + 1)^{-1}/4 + \|x_1\|\|x_2\| \leq \|x_1\|\|x_2\| + 1.$$

But by the way we have chosen K , x_{12} and x_{21} have disjoint supports, hence $x_{12} \perp x_{21}$. Thus

$$\begin{aligned} & |\langle ux_1 u^* x_2, x_2 u x_1 u^* \rangle| \\ & \leq |\langle ux_1 u^* x_2 - x_{12}, x_2 u x_1 u^* \rangle| + |\langle x_{12}, x_2 u x_1 u^* - x_{21} \rangle| \\ & \leq \|ux_1 u^* x_2 - x_{12}\|_2 \|x_2 u x_1 u^*\|_2 + \|x_{12}\|_2 \|x_2 u x_1 u^* - x_{21}\|_2 \\ & \leq \varepsilon(\|x_1\|\|x_2\| + 1)^{-1} \|x_1\|\|x_2\|/4 + (\|x_1\|\|x_2\| + 1)(\varepsilon(\|x_1\|\|x_2\| + 1)^{-1}/4) < \varepsilon. \end{aligned}$$

□

1.4.4. Proposition. *With the same notations and assumptions as in 1.4.2 and 1.4.3, let $q \in \mathcal{P}(B)$ and let $A_0 \subset qMq$ be a diffuse abelian von Neumann subalgebra. Assume no corner of $(\mathcal{N}_{qMq}(A_0) \cap qM_1q)'' \subset qMq$ can be embedded into B inside M_1 . Then a corner of A_0 can be embedded into qM_1q inside qMq .*

Proof. Assume that no corner of A_0 embeds into M_1 . Apply first 1.4.2 for $\varepsilon = \tau(q)^{1/2}/4$ to deduce that there exists $F \subset \Lambda \setminus \Lambda_1$ finite and $x_1, x_2 \in M$ supported on F such that if $u \in \mathcal{N}_{qMq}(A_0)$ then $\|ux_1 u^* x_2 - x_2 u x_1 u^*\|_2 \leq \tau(q)^{1/2}/4$ and $3\tau(q)^{1/2}/4 \leq \|ux_1 u^* x_2\|_2 \leq 5\tau(q)^{1/2}/4$. It then follows that $|\langle ux_1 u^* x_2, x_2 u x_1 u^* \rangle| \geq \tau(q)/4, \forall u \in \mathcal{N}_{qMq}(A)$.

By Lemma 1.4.3, there exist $K \subset \Lambda_1$ finite and $\delta > 0$ such that if $u \in (M_1)_1$ satisfies $\|E_B(uu_g^*)\|_2 \leq \delta, \forall g \in K$, then $ux_1 u^* x_2 \perp_{\tau(q)/4} x_2 u x_1 u^*$. But this implies that we cannot find $u \in \mathcal{N} = \mathcal{N}_{qMq}(A_0) \cap qM_1q$ such that $\|E_B(uu_g^*)\|_2 \leq \delta, \forall g \in K$. By (2.3 in [P6]) this contradicts the fact that no corner of $\mathcal{N}''$ embeds into B inside M_1 . □

The next result provides a useful “transitivity” property for the subordination relation considered in Corollary 1.2.4:

1.4.5. Lemma. *Let M be a finite von Neumann algebra, $B_0, M_1 \subset M$ von Neumann subalgebras of M and $Q \subset M_1$ a von Neumann subalgebra of M_1 . Assume there exist projections $q_0 \in B_0$, $q_1 \in M_1$, a unital isomorphism of $q_0 B_0 q_0$ into $q_1 M_1 q_1$ and a partial isometry $v \in M$ such that $v^* v \in (q_0 B_0 q_0)' \cap q_0 M q_0$, $vv^* \in \psi(q_0 B_0 q_0)' \cap q_1 M q_1$ and $vb = \psi(b)v, \forall b \in q_0 B_0 q_0$. Denote by q' the support projection of $E_{M_1}(vv^*) \in \psi(q_0 B_0 q_0)' \cap q_1 M_1 q_1$ and $B_1 = \psi(q_0 B_0 q_0)q'$. If a corner of $B_1 = \psi(q_0 B_0 q_0)q'$ can be embedded into Q inside M_1 , then a corner of B_0 can be embedded into Q inside M .*

Proof. Indeed, if $p_1 \in \mathcal{P}(B_1)$, $v_1 \in M_1 p_1$ is a non-zero partial isometry and $\psi_1 : p_1 B_1 p_1 \rightarrow Q$ a (not necessarily unital) isomorphism such that $v_1 b = \psi_1(b)v_1, \forall b \in p_1 B_1 p_1$, then $v_1 v \neq 0$ and $v_1 v b = \psi_1(\psi(b))v_1 v, \forall b \in q_0 B_0 q_0$. $\square$

Proof of Theorem 1.4.1. By Proposition 1.4.4 and (2.1 in [P6]), there exist projections $q_0 \in A_0 \subset qMq$, $q_1 \in qM_1q$, a unital isomorphism of $A_0 q_0$ into $q_1 M_1 q_1$ and a partial isometry $v \in M$ such that $v^* v = q_0$, $vv^* \in \psi(A_0 q_0)' \cap q_1 M q_1$ and $va = \psi(a)v, \forall a \in A_0 q_0$. Let q' be the support projection of $E_{M_1}(vv^*)$ and note that if we denote $A_1 = \psi(A_0 q_0) \subset q_1 M_1 q_1$ then $q' \in A_1' \cap q_1 M_1 q_1$. By replacing if necessary ψ by $q'\psi(\cdot)q'$ and shrinking $q_0 \in A_0$ accordingly, we may assume $q_1 = q'$.

Now, if a corner of A_1 can be embedded into A inside M_1 , then by Lemma 1.4.5 a corner of A_0 can be embedded into A inside M , so by (A.1 in [P5]) the two Cartan subalgebras $A_0, Aq \subset qMq$ are unitary conjugate. If in turn no corner of A_1 can be embedded into A inside M_1 then by Theorem 1.2.1 we have $vv^* \in A_1' \cap q_1 M q_1 \subset q_1 M_1 q_1$, implying that $vA_0 v^* \subset q_1 M_1 q_1$. By spatiality, $vA_0 v^*$ is Cartan in $q_1 M q_1$, which by Corollary 1.2.4 implies that a corner of $vA_0 v^*$ can be embedded into A inside M_1 . By (A.1 in [P5]), this implies A_0 and Aq are unitary conjugate in qMq . On the other hand, by Proposition 1.3.1 we have $A_0 \subset qM_1 q$. But then Theorem 1.2.1 applies to show that A_0, Aq are conjugate in $qM_1 q$ as well. $\square$

2. DEFORMATION OF AFP FACTORS.

Throughout this section, $(M_1, \tau_1), (M_2, \tau_2)$ will be finite von Neumann algebras with $B \subset M_i$ a common von Neumann subalgebra such that $\tau_1|_B = \tau_2|_B$, as in Section 1. We describe in this section several useful ways to deform the identity map on the AFP algebra $M = M_1 *_B M_2$ by subunital, subtracial, c.p. maps which arise naturally from the amalgamated free product structure of M . By a *deformation* of id_M we mean here a sequence ϕ_n of subunital, subtracial, c.p. maps on M such that $\lim_n \|\phi_n(x) - x\|_2 = 0, \forall x \in M$.

2.1. Amalgamated free product of c.p. maps. We first recall the definition of amalgamated product of c.p. maps from ([Bo]), and establish some basic properties.

2.1.1. Lemma. *Let $\phi_i : M_i \rightarrow M_i$ be subunital, subtracial, c.p. maps with $\phi_1(b) = \phi_2(b), \forall b \in B$, and $\tau \circ \phi_i = \lambda \tau, \forall i = 1, 2$, with $0 < \lambda \leq 1$. Then $\phi_1 *_B \phi_2 : M_1 *_B M_2 \rightarrow M_1 *_B M_2$ is a well-defined subunital, subtracial, c.p. map. Moreover, the map $(\phi_1, \phi_2) \mapsto \phi_1 *_B \phi_2$ is continuous with respect to the topologies given by pointwise $\|\cdot\|_2$ -convergence.*

Proof. Since $\tau \circ \phi_i = \lambda\tau$, $\forall i = 1, 2$ we have that $\tau \circ (\phi_1 *_B^{\text{alg}} \phi_2) = \lambda\tau$, and hence by [Bo] extends uniquely to a c.p. map $\phi_1 *_B \phi_2$ on $M_1 *_B M_2$ which follows subtracial.

To show that this correspondence is continuous suppose that $\varepsilon > 0$ and $x_1, \dots, x_n \in M_1 *_B M_2$. As $M_1 *_B^{\text{alg}} M_2$ is dense in $M_1 *_B M_2$ let $x'_1, \dots, x'_n \in M_1 *_B^{\text{alg}} M_2$ such that $\|x_j - x'_j\|_2 \leq \varepsilon/3$, $\forall j \leq n$. Hence $\exists m, l \in \mathbb{N}$, $F_i \subset M_i$ finite, such that each x'_i is the sum of at most m products of length at most l from $F_1 \cup F_2$. Let $N = \max_{x \in F_1 \cup F_2} \|x\|$, then if $\phi'_i : M_i \rightarrow M_i$ are subunital, subtracial, c.p. maps with $\|\phi'_i(x) - \phi_i(x)\|_2 < \varepsilon/3nlN^l$, $\forall x \in F_i$, $i = 1, 2$, then repeated use of the triangle inequality together with the fact that subunital c.p. maps are contractions in the uniform norm show that $\|\phi'_1 *_B \phi'_2(x_i) - \phi_1 *_B \phi_2(x_i)\|_2 < \varepsilon$, $\forall i \leq n$. $\square$

2.1.2. Remark. In general, the free product of two subunital subtracial c.p. maps need not be subtracial. In fact, given any c.p. map ϕ_1 which is unital, subtracial, but not tracial, the c.p. map $\phi = \phi_1 * id$ is not subtracial. Even more so, the Radon-Nykodim derivative $d\tau \circ \phi/d\tau$ of any such free product c.p. map ϕ is unbounded. To see this, let $x \in M_1$ such that $\tau(x) = 0$, but $\tau \circ \phi_1(x) \neq 0$, let $v \in M_2$ be a partial isometry with $\tau(v) = 0$ and set $p = vv^*$. Then we have:

$$\begin{aligned} \tau \circ \phi((v xv^*)^*(v xv^*)) &= |\tau \circ \phi_1(x)|^2 \tau(p) + \|v(\phi_1(x) - \tau \circ \phi_1(x))v^*\|_2^2 \\ &= |\tau \circ \phi_1(x)|^2 \tau(p) + \tau(p)^2 \|\phi_1(x) - \tau \circ \phi_1(x)\|_2^2. \end{aligned}$$

Thus, if we choose v such that $\tau(p) \rightarrow 0$ then

$$\tau \circ \phi((v xv^*)^*(v xv^*)) / \tau((v xv^*)^*(v xv^*)) \rightarrow \infty.$$

We note that in [Bo] it was assumed that a free product of subtracial c.p. maps is subtracial in order to show that the Haagerup property is preserved by free products. But although the above argument shows that this fact doesn't hold true unless the c.p. maps are actually tracial, the result on the Haagerup property is still valid, since by (Proposition 2.2. in [Jol]) one can take the c.p. maps given by the Haagerup property to be unital and tracial.

2.2. Deformation by automorphisms. Suppose $\alpha_i \in \text{Aut}(M_i, \tau_{M_i})$ such that $\alpha_1(b) = \alpha_2(b)$, $\forall b \in B$. Then since automorphisms are unital c.p. maps we have that $\alpha = \alpha_1 *_B \alpha_2$ is a unital, tracial, c.p. map, moreover α restricted to the dense subalgebra $M_1 *_B^{\text{alg}} M_2$ is an automorphism and so by continuity we have that α is an automorphism.

Hence if $\alpha_i^t \in \text{Aut}(M_i)$, $t \in \mathbb{R}$ is a one parameter group of automorphisms of M_i which is pointwise $\|\cdot\|_2$ -continuous and satisfies $\alpha_{i|B}^t = id_B$, for $i = 1, 2$, then α^t gives a deformation of the identity of M by automorphisms. In particular, we have:

2.2.1. Lemma. *Let $v_i \in \mathcal{U}(B' \cap M_i)$, $i = 1, 2$. Then there is a pointwise $\|\cdot\|_2$ -continuous one parameter group of automorphisms $\{\alpha^t\}_{t \in \mathbb{R}} \subset \text{Aut}(M)$ such that $\alpha_1 = \text{Ad}(v_1) *_B \text{Ad}(v_2)$.*

Proof. Let $h_j = h_j^* \in B' \cap M_j$ be such that $\exp(\pi i h_j) = v_j$ and define $\alpha_j^t = \text{Ad}(\exp(\pi t i h_j))$, $t \in \mathbb{R}$, $j = 1, 2$, and the above observation applies. $\square$

For the next lemma we denote $\tilde{M} = M *_B (B \overline{\otimes} L(\mathbb{F}_2))$. Note that if we let $L(\mathbb{F}_2) = L(\mathbb{Z} * \mathbb{Z}) = L(\mathbb{Z}) * L(\mathbb{Z})$ and denote $\tilde{M}_i = M_i *_B (B \overline{\otimes} L(\mathbb{Z}))$, $i = 1, 2$, then $\tilde{M} = \tilde{M}_1 *_B \tilde{M}_2$. Also, if $u_1 \in L(\mathbb{Z} * 1) \subset L(\mathbb{F}_2)$, $u_2 \in L(1 * \mathbb{Z}) \subset L(\mathbb{F}_2)$ are the canonical generating unitaries then $u_i \in B' \cap \tilde{M}_i$, $i = 1, 2$. We will use the algebra $\tilde{M}$ as framework for the main deformation of M , 2.2.2 below. The action of M on the (1.1.1)-decomposition of the AFP algebra $\tilde{M} = \tilde{M}_1 *_B \tilde{M}_2$ can be viewed as the analogue of the action of an amalgamated free product group $\Lambda_1 *_H \Lambda_2$ on the Bass-Serre tree with vertices $\Lambda_1/H \cup \Lambda_2/H$.

In turn, the “graded deformation” below is inspired by the graded deformations of crossed product algebras involving Bernoulli actions in ([P3], [P6]).

2.2.2. Lemma. *There exists a pointwise $\|\cdot\|_2$ -continuous one parameter group of automorphisms $\{\theta_t\}_{t \in \mathbb{R}}$ and a period 2 automorphism β of $\tilde{M}$ such that:*

- (a). $\theta_0 = \text{id}$, $\theta_1 = \text{Ad}(u_1) *_B \text{Ad}(u_2)$.
- (b). $\beta \theta_t \beta = \theta_{-t}$, $\forall t \in \mathbb{R}$.
- (c). $M \subset \tilde{M}^\beta$.

Proof. Let A_j be the von Neumann subalgebra generated by u_j , and let $h_j \in A_j$ be self-adjoint elements with spectrum in $[-\pi, \pi]$ such that $u_j = \exp(\pi i h_j)$. Set $u_j^t = \exp(\pi t i h_j)$, $j = 1, 2$, $t \in \mathbb{R}$. Then $\theta_t = \text{Ad}(u_1^t) *_B \text{Ad}(u_2^t) \in \text{Aut}(\tilde{M})$, $\forall t \in \mathbb{R}$, defines a pointwise $\|\cdot\|_2$ -continuous one parameter group of automorphisms which satisfies (a).

Let β be the unique automorphism of $\tilde{M}$ satisfying $\beta|_M = \text{id}_M$ and $\beta(u_j) = u_j^*$, $j = 1, 2$. Then β is clearly a period 2 automorphism and it satisfies (c) by definition. Also, for $x \in M = M_1 *_B M_2$ we have

$$\begin{aligned} \beta \theta_t \beta(x) &= \beta \theta_t(x) = \beta(\text{Ad}(\exp(\pi t i h_1)) * \text{Ad}(\exp(\pi t i h_2)))(x) \\ &= (\text{Ad}(\exp(-\pi t i h_1)) * \text{Ad}(\exp(-\pi t i h_2)))(x) = \theta_{-t}(x), \forall x \in M. \end{aligned}$$

Similarly, for u_1, u_2 we have

$$\beta \theta_t \beta(u_j) = \beta \theta_t(u_j^*) = \beta(u_j^*) = u_j = \theta_{-t}(u_j).$$

Since u_1, u_2 , and M generate $\tilde{M}$ as a von Neumann algebra it follows that $\beta \theta_t \beta = \theta_{-t}$, $\forall t$. $\square$

2.3. Deformation by free products of multiples of the identity. Recall that if $\mathcal{H}_i^0 = L^2(M_i) \ominus L^2(B)$ then we may decompose $L^2(M_1 *_B M_2)$ in the usual way as

$$L^2(M_1 *_B M_2) = L^2(B) \oplus \bigoplus_{n \geq 1} \bigoplus_{\substack{i_j \in \{1, 2\} \\ i_1 \neq i_2, \dots, i_{n-1} \neq i_n}} \mathcal{H}_{i_1}^0 \otimes_B \mathcal{H}_{i_2}^0 \otimes_B \cdots \otimes_B \mathcal{H}_{i_n}^0.$$

For each $L \in \mathbb{N}$ we let $\widehat{E}_L$ be the projection onto the subspace

$$\bigoplus_{n \geq L} \bigoplus_{\substack{i_j \in \{1,2\} \\ i_1 \neq i_2, \dots, i_{n-1} \neq i_n}} \mathcal{H}_{i_1}^0 \otimes_B \mathcal{H}_{i_2}^0 \otimes_B \cdots \otimes_B \mathcal{H}_{i_n}^0.$$

Let $\{c_n\}_n \subset [0, 1)$ such that $c_n \nearrow 1$, then by Lemma 1.1.1 we have:

2.3.1. Lemma. $\phi_n = (c_n id) *_B (c_n id)$ gives a deformation of the identity of M . Moreover, ϕ_n commutes with $\widehat{E}_L$ as operators on $L^2(M)$ and $\|\phi_n \circ \widehat{E}_L(x)\|_2 \leq c_n^L \|x\|_2$, $\forall n, L \in \mathbb{N}$, $x \in M$.

Proof. Trivial by the definitions.

2.4. Deformation by subalgebras. For each $i \in \{1, 2\}$ let $N_i^j \subset M_i$ be an increasing sequence of von Neumann subalgebras such that $B \subset N_i^1$, and $\overline{\bigcup_{j \geq 1} N_i^j} = M_i$. Let $E_i^j : M_i \rightarrow M_i$ be the conditional expectation onto N_i^j . Then $E^j = E_1^j *_B E_2^j$ gives a sequence of conditional expectations of M onto $N_1^j *_B N_2^j$, which by 2.1.1 converges to the identity pointwise, i.e. $\overline{\bigcup_{j \geq 1} N_1^j *_B N_2^j} = M$.

A particular case of such a deformation, which works whenever $B' \cap M_i$ is diffuse, is given by $N_i^j = p_i^j M_i p_i^j \oplus B(1 - p_i^j)$, with $p_i^j \in \mathcal{P}(B' \cap M_i)$ satisfying $p_i^j \nearrow 1_{M_i}$, $i = 1, 2$. Indeed, this clearly implies $\overline{\bigcup_{j \geq 1} (p_1^j M_1 p_1^j \oplus B(1 - p_1^j)) *_B (p_2^j M_2 p_2^j \oplus B(1 - p_2^j))} = M$.

3. DEFORMATION/RIGIDITY ARGUMENTS.

In this section we investigate the effect that the deformations considered in Section 2 have on the relatively rigid subalgebras of $M_1 *_B M_2$. To this end, first recall from (Sec. 4 in [P5]) that if $Q \subset M$ is a von Neumann subalgebra of the finite von Neumann algebra (M, τ) , then $Q \subset M$ is called a *rigid inclusion* (or Q is a *relatively rigid subalgebra* of M) if any deformation of id_M by subunital subtracial c.p. maps $\{\phi_n\}_n$ tends uniformly to id_Q on the unit ball of Q , i.e. $\lim_n \sup\{\|\phi_n(y) - y\|_2 \mid y \in Q, \|y\| \leq 1\} = 0$. This property doesn't in fact depend on the choice of the trace τ on M and can be given several other equivalent characterizations (see [P5], [PeP]). The following provides yet another characterization of relative rigidity, by showing that it is enough to consider deformations by unital tracial c.p. maps:

3.1. Theorem. *Let N be a finite von Neumann algebra with countable decomposable center. Let Q be a von Neumann subalgebra, then the following are equivalent:*

- (i). *The inclusion $Q \subset N$ is rigid.*
- (ii). *There exists a normal faithful tracial state τ on N such that: $\forall \varepsilon > 0$, $\exists F = F(\varepsilon) \subset N$ finite and $\delta = \delta(\varepsilon) > 0$ such that if $\phi : N \rightarrow N$ is a normal, c.p. map with $\tau \circ \phi = \tau$, $\phi(1) = 1$, and $\|\phi(x) - x\|_2 \leq \delta$, $\forall x \in F$, then $\|\phi(b) - b\|_2 \leq \varepsilon$, $\forall b \in Q$, $\|b\| \leq 1$.*
- (iii). *Condition (ii) is satisfied for any normal faithful tracial state τ on N .*

Proof. (i) $\implies$ (iii) and (iii) $\implies$ (ii) are both trivial and so it is enough to show (ii) $\implies$ (i). That is, assuming (ii) holds we must show that the following condition holds: (3.1.1) $\forall \varepsilon > 0$, $\exists F' = F'(\varepsilon) \subset N$ finite and $\delta' = \delta'(\varepsilon) > 0$ such that if $\phi : N \rightarrow N$ is a normal, c.p. map with $\tau \circ \phi \leq \tau$, $\phi(1) \leq 1$, and $\|\phi(x) - x\|_2 \leq \delta'$, $\forall x \in F'$, then $\|\phi(b) - b\|_2 \leq \varepsilon$, $\forall b \in Q$, $\|b\| \leq 1$.

By (Lemma 3 in [PeP]) we may also assume that in the above condition ϕ is symmetric, i.e. $\tau(\phi(x)y) = \tau(x\phi(y))$, $\forall x, y \in N$. Let $F = F(\varepsilon/2)$, and $\delta = \delta(\varepsilon/2)$ be given from (ii). Let $F' = F \cup \{1\}$, and $\delta' = \min_{x \in F} \{\delta/2, \delta/(8\|x\|^2 + 1), \varepsilon^2/8\}$, suppose that $\phi : N \rightarrow N$ is a normal, symmetric c.p. map with $\tau \circ \phi \leq \tau$, $\phi(1) \leq 1$, and $\|\phi(x) - x\|_2 \leq \delta'$, $\forall x \in F'$. Let $a = \phi(1) = d\tau \circ \phi/d\tau$ and define ϕ' by $\phi'(x) = \phi(x) + (1-a)^{1/2}x(1-a)^{1/2}$. Then ϕ' is a normal, c.p. map with $\phi'(1) = 1$. Moreover, as ϕ is symmetric, so is ϕ' and hence $\tau \circ \phi' = \tau$.

Also, it follows that for each $x \in F$ we have

$$\begin{aligned} \|\phi'(x) - x\|_2 &\leq \|\phi(x) - x\|_2 + \|(1-a)^{1/2}x(1-a)^{1/2}\|_2 \\ &\leq \|\phi(x) - x\|_2 + \|(1-a)x(1-a)\|_2^{1/2}\|x\|^{1/2} \\ &\leq \|\phi(x) - x\|_2 + \|1-a\|_2\|x\| \leq \delta. \end{aligned}$$

Hence, by (ii) we have $\|\phi'(b) - b\|_2 \leq \varepsilon/2$, $\forall b \in Q$, $\|b\| \leq 1$. Thus

$$\begin{aligned} \|\phi(b) - b\|_2 &\leq \|\phi'(b) - b\|_2 + \|(1-a)^{1/2}x(1-a)^{1/2}\|_2 \\ &\leq \|\phi'(b) - b\|_2 + \|1-a\|_2 \leq \varepsilon, \end{aligned}$$

for all $b \in Q$ with $\|b\| \leq 1$. □

3.2. Corollary. *Let (M_1, τ_1) and (M_2, τ_2) be finite von Neumann algebras and $Q \subset M_1$ a von Neumann subalgebra such that the inclusion $Q \subset M_1 * M_2$ is rigid. Then the inclusion $Q \subset M_1$ is rigid.*

Proof. Let $\varepsilon > 0$ be given. Since $Q \subset M_1 * M_2$ is rigid we can find $F = F(\varepsilon) \subset M$ finite and $\delta = \delta(\varepsilon) > 0$ to satisfy condition (ii) of Theorem 3.1. By Lemma 2.1.1 there exists $F' \subset M_1$ finite, and $\delta' > 0$ such that if $\phi : M_1 \rightarrow M_1$ is a normal unital, tracial c.p. map such that $\|\phi(x) - x\|_2 \leq \delta'$, $\forall x \in F'$ then $\|\phi * id_{M_2}(x) - x\|_2 \leq \delta$, $\forall x \in F$. Hence by our choice of F and δ we have that $\|\phi(b) - b\|_2 = \|\phi * id_{M_2}(b) - b\|_2 \leq \varepsilon$, $\forall b \in Q$, $\|b\| \leq 1$. Thus, by 3.1 $Q \subset M_1$ follows a rigid inclusion. □

We'll now use the deformation 2.2.2 to exploit the relative rigidity of subalgebras $Q \subset M_1 *_B M_2 \subset \tilde{M}_1 *_B \tilde{M}_2$. This “deformation/rigidity” argument is inspired from (4.3-4.8 in [P3] and Sec. 4 in [P6I])

3.3. Proposition. *Let $(M_1, \tau_1), (M_2, \tau_2)$ be finite von Neumann algebras with a common von Neumann subalgebra $B \subset M_i, i = 1, 2$, such that $\tau_{1|B} = \tau_{2|B}$. Denote $M = M_1 *_B M_2$, $\tilde{M}_i = M_i *_B (B \overline{\otimes} L(\mathbb{Z}))$, $i = 1, 2$, and $\tilde{M} = \tilde{M}_1 *_B \tilde{M}_2 = M *_B (B \overline{\otimes} L(\mathbb{F}_2))$, as in 2.2.2.*

Denote $\theta = \text{Ad}(u_1) * \text{Ad}(u_2) \in \text{Aut}(\tilde{M})$, where $u_1, u_2 \in L(\mathbb{F}_2)$ are the canonical generators of $L(\mathbb{F}_2)$, as in 2.2.2. Let $Q \subset M$ be a von Neumann subalgebra such that $Q \subset \tilde{M}$ is a rigid inclusion and assume no corner of Q can be embedded into B inside M , i.e. $Q' \cap \langle M, B \rangle$ contains no non-zero finite projections. Then there exists a non-zero partial isometry $v \in \tilde{M}$ such that $vy = \theta(y)v, \forall y \in Q$.

Proof. Since $Q \subset \tilde{M}$ is rigid, there exist $F \subset \tilde{M}$ finite and $\delta > 0$ such that if $\phi : \tilde{M} \rightarrow \tilde{M}$ is a subunital, subtracial, c.p. map with $\|\phi(x) - x\|_2 \leq \delta, \forall x \in F$, then $\|\phi(u) - u\|_2 \leq 1/2, \forall u \in \mathcal{U}(Q)$. Using the continuity of $t \mapsto \theta_t$, we can find $n \geq 1$ such that $\|\theta_{1/2^n}(x) - x\| \leq \delta, \forall x \in F$, which entails $\|\theta_{1/2^n}(u) - u\| \leq 1/2, \forall u \in \mathcal{U}(Q)$.

Now, let a be the unique element of minimal norm- $\|\cdot\|_2$ in $K = \overline{\text{co}}^w\{\theta_{1/2^n}(u)u^* \mid u \in \mathcal{U}(Q)\}$. From $\|\theta_{1/2^n}(u)u^* - 1\|_2 \leq 1/2, \forall u \in \mathcal{U}(Q)$, we get that $\|a - 1\|_2 \leq 1/2$, so $a \neq 0$. Since $\theta_{1/2^n}(u)Ku^* = K$ and $\|\theta_{1/2^n}(u)au^*\|_2 = \|a\|_2, \forall u \in \mathcal{U}(Q)$ we deduce, using the uniqueness of a , that $au = \theta_{1/2^n}(u)a, \forall u \in \mathcal{U}(Q)$. Using standard arguments we can replace a by the partial isometry in its polar decomposition, thus getting a non-zero partial isometry $v \in M$ such that $vu = \theta_{1/2^n}(u)v, \forall u \in \mathcal{U}(Q)$.

In what follows we show by induction that for any $k \geq 0$ there exist a non-zero partial isometry $v_k \in \tilde{M}$ such that

$$(3.3.1) \quad v_k u = \theta_{1/2^{n-k}}(u)v_k, \forall u \in \mathcal{U}(Q),$$

which for $k = n$ gives the conclusion. Since we have already constructed v_0 we only need to construct v_{k+1} , given v_k . Note first that if v_k satisfies (3.3.1) then $v_k^* v_k \in Q' \cap \tilde{M}$ and $v_k v_k^* \in \theta_{1/2^{n-k}}(Q)' \cap \tilde{M}$.

By applying the automorphism β to the equality (3.3.1) and using the properties of β we get:

$$(3.3.2) \quad \begin{aligned} \beta(v_k)u &= \beta(v_k)\beta(u) = \beta(\theta_{1/2^{n-k}}(u))\beta(v_k) \\ &= \theta_{-1/2^{n-k}}(\beta(u))\beta(v_k) = \theta_{-1/2^{n-k}}(u)\beta(v_k), \forall u \in \mathcal{U}(Q). \end{aligned}$$

By replacing u by u^* and taking conjugates in (3.3.2), we obtain

$$u\beta(v_k^*) = \beta(v_k^*)\theta_{-1/2^{n-k}}(u), \forall u \in \mathcal{U}(Q),$$

which combined with (3.1.1) gives

$$(3.3.3) \quad \begin{aligned} v_k \beta(v_k^*) \theta_{-1/2^{n-k}}(u) &= v_k u \beta(v_k^*) \\ &= \theta_{1/2^{n-k}}(u) v_k \beta(v_k^*), \forall u \in \mathcal{U}(Q). \end{aligned}$$

By applying $\theta_{1/2^{n-k}}$ to (3.3.3), we further get for $u \in \mathcal{U}(Q)$ the identity

$$(3.3.4) \quad \theta_{1/2^{n-k}}(v_k \beta(v_k^*))u = \theta_{1/2^{n-k-1}}(u) \theta_{1/2^{n-k}}(v_k \beta(v_k^*)).$$

Since no corner of Q can be embedded into B inside M we can apply Theorem 1.1 to conclude that $Q' \cap \tilde{M} \subset M$. Thus, since $v_k^* v_k \in Q' \cap \tilde{M}$ and $M \subset \tilde{M}^\beta$, we get $\beta(v_k^* v_k) = v_k^* v_k$, implying that $v_k \beta(v_k^*)$ is a partial isometry of same left support as v_k . Thus, by (3.3.4), $w = \theta_{1/2^{n-k}}(v_k \beta(v_k^*))$ is a non-zero partial isometry satisfying $wu = \theta_{1/2^{n-k-1}}(u)w, \forall u \in \mathcal{U}(Q)$, and the inductive step follows. $\square$

3.4. Proposition. *As in 2.3, denote by $\widehat{E}_L$ the orthogonal projection of $L^2(M)$ onto the Hilbert space spanned by reduced words of length $\geq L$. If $Q \subset M_1 *_B M_2$ is a rigid inclusion, then for any $\varepsilon > 0$, there exists $L \in \mathbb{N}$ such that $\|\widehat{E}_L(x)\|_2 < \varepsilon$, $\forall x \in Q$, $\|x\| \leq 1$.*

Proof. Let $\phi_n : M \rightarrow M$ be as in 2.3.1, for some $c_n \nearrow 1$, then by 2.1 we have $\lim_n \|\phi_n(x) - x\|_2 = 0$, $\forall x \in M$. Thus, since $Q \subset M$ is rigid, there exists $l \in \mathbb{N}$ such that $\|\phi_l(x) - x\|_2 < \varepsilon/2$, $\forall x \in Q$, $\|x\| \leq 1$. Let $L \in \mathbb{N}$ such that $c_l^L < \varepsilon/2$. Then

$$\begin{aligned} \|\widehat{E}_L(x)\|_2 &\leq \|\widehat{E}_L(x - \phi_l(x))\|_2 + \|\phi_l \circ \widehat{E}_L(x)\|_2 \\ &\leq \|x - \phi_l(x)\|_2 + c_l^L \|x\|_2 < \varepsilon, \forall x \in Q, \|x\| \leq 1. \end{aligned}$$

□

4. EXISTENCE OF INTERTWINING BIMODULES.

In the previous section we saw that a relatively rigid subalgebra Q of a finite AFP von Neumann algebra $M = M_1 *_B M_2$ can be “located” by certain c.p. deformations of id_M . In this section we will use this information to prove that $L^2(M)$ must contain non-trivial Hilbert bimodules intertwining Q into either M_1 or M_2 . The rather long and technical proof will proceed by contradiction, assuming Q cannot be intertwined in neither M_1 nor M_2 , inside M . We first show that this implies Q cannot be intertwined into M_1, M_2 in $\tilde{M} = (M_1 *_B M_2) *_B (B \overline{\otimes} L(\mathbb{F}_2))$ either. By Proposition 3.4 and (2.3 in [P6]), this shows that Q must contain “at infinity” elements with uniformly bounded free length and at least two “very large letters” in M_1, M_2 or $L(\mathbb{F}_2)$. This will be shown to contradict Proposition 3.3.

To “measure” the letters in M_i we will need these algebras to have nice orthonormal basis over B , in the following sense:

4.1. Definition. Let (M, τ) be separable finite von Neumann algebra and $B \subset M$ a von Neumann subalgebra. A sequence of elements $\{\eta_n\}_n \subset M$ satisfying the conditions $\eta_0 = 1$, $E_B(\eta_i^* \eta_j) = \delta_{ij}$, $\forall i, j$ and $\Sigma \eta_n B$ dense in $L^2(M, \tau)$ is called a *bounded, homogeneous, orthonormal basis* (BHOB) of M over B . An inclusion $B \subset M$ having a BHOB is said to be *homogeneous*.

4.2. Lemma. *Let (M, τ) be a separable finite von Neumann algebra and $B \subset M$ a von Neumann subalgebra. Assume one of the following conditions holds true:*

- (a). $B = \mathbb{C}1$.
- (b). $B = A \subset M$ is Cartan (i.e. B is maximal abelian and regular in M) and M is type II_1 .
- (c). $B = N \subset M$ is an irreducible inclusion of II_1 factors and B is regular in M .

Then $B \subset M$ is homogeneous, moreover in both cases (b) and (c), M has a BHOB made of unitary elements in $\mathcal{N}_M(B)$.

Proof. Case (a) is clear by the Gram-Schmidt algorithm. The case (c) is trivial once we notice that such $N \subset M$ is a crossed product inclusion $N \subset M = N \rtimes_{\sigma, v} G$ for some cocycle action σ of a discrete countable group G on N , with 2-cocycle v . Indeed, in this case the canonical unitaries $\{u_g\}_g \subset M$ implementing the action σ provide a BHOB of M over N .

To prove the case (b), we first show that given any $n \geq 1$ and any finite set $v_0 = 1, v_1, \dots, v_{n-1} \in \mathcal{N}_M(A)$, with $E_A(v_i^* v_j) = 0$ for $0 \leq i \neq j < n$, and $v'_n \in \mathcal{GN}_M(A)$, with $v'_n{}^* v'_n \neq 1$, there exists $v \in \mathcal{GN}_M(A)$ non-zero such that $v^* v'_n = 0, v'_n v^* = 0$ and $E_A(v^* v_j) = 0, \forall 1 \leq j \leq n-1$. Indeed, note first that by ([Dy]) there exists $u \in \mathcal{N}_M(A)$ such that $v'_n = u(1-p)$, where $p = 1 - v'_n{}^* v'_n \in \mathcal{P}(A)$. Since $Ap \subset pMp$ is Cartan with pMp of type II_1 , it follows that there exists $w \in \mathcal{N}_{pMp}(Ap)$ such that $w \notin (\Sigma_{i=0}^{n-1} pu^* v_i Ap)$ (or else, pMp would follow finite dimensional over Ap thus of type I, implying that M has a type I direct summand, a contradiction). Then $w_i = pu^* v_i p \in \mathcal{GN}(Ap)$, $0 \leq i \leq n-1$, satisfy $0 \neq w - \Sigma_{i=0}^{n-1} w_i E_A(w_i^* w) \in \mathcal{GN}(A)$ (by [Dy]). But then $v = u(w - \Sigma_{i=0}^{n-1} w_i E_A(w_i^* w))$ clearly satisfies all required conditions.

Now, to finish the proof of (b) let $\{u_n\}_{n \geq 0} \subset \mathcal{N}_M(A)$ be a sequence of unitaries normalizing A , dense in $\mathcal{N}_M(A)$ in the norm $\|\cdot\|_2$, with $u_0 = 1$ and with each u_n appearing with infinite multiplicity. It is sufficient to construct a sequence $\{v_n\}_{n \geq 0} \subset \mathcal{N}_M(A)$ such that for all $m \geq 0$ we have:

$$(4.2.1) \quad E_A(v_i^* v_j) = \delta_{ij}, \forall m \geq i, j \geq 0, u_m \in \Sigma_{i=0}^m v_i A.$$

Assume we have constructed $v_0 = 1, v_1, \dots, v_{n-1}$ satisfying (4.2.1) for $m = n-1$, for some $n \geq 1$. Let $v''_n = u_n - \Sigma_{i=0}^{n-1} v_i E_A(v_i^* u_n) \in \mathcal{GN}_M(A)$. Let $v'_n \in \mathcal{GN}_M(A)$ be maximal with the properties: $v'_n (v''_n)^* v''_n = v''_n$ and $v'_n \perp \Sigma_{i=0}^{n-1} v_i A$. By applying the first part of the proof to $v_0, v_1, \dots, v_{n-1}, v'_n$ and using the maximality, it follows that v'_n is a unitary element. But then $v_n = v'_n$ clearly satisfies (4.2.1). $\square$

4.3. Theorem. *Let (M_1, τ_1) and (M_2, τ_2) be finite von Neumann algebras and $B \subset M_i$ a common von Neumann subalgebra such that $\tau_1|_B = \tau_2|_B$. Suppose that the inclusions $B \subset M_1$ and $B \subset M_2$ are homogeneous. Let Q be a von Neumann subalgebra of $M = M_1 *_B M_2$ such that the inclusion $Q \subset M$ is rigid. Then for either $i = 1$ or $i = 2$ there exists a non-zero projection f in $Q' \cap \langle M, e_{M_i} \rangle$ of finite trace $Tr = Tr_{\langle M, e_{M_i} \rangle}$.*

Proof. By taking spectral projections, it is sufficient to show there exists $i \in \{1, 2\}$ and $a \in Q' \cap \langle M, e_{M_i} \rangle$ with $0 \leq a \leq 1, 0 \neq Tr(a) < \infty$. Assume by contradiction that there are no such elements in $Q' \cap \langle M, e_{M_i} \rangle, i = 1, 2$. If we identify Q with the diagonal subalgebra $\{x \oplus x \mid x \in Q\}$ in $M \oplus M$, then this is equivalent to saying that Q cannot be intertwined into $M_1 \oplus M_2$ inside $M \oplus M$. By (2.3 in [P6]) this implies the following:

Fact 1: $\forall \varepsilon > 0, \forall y_1, \dots, y_n \in M, \exists w \in \mathcal{U}(Q)$ such that $\|E_{M_i}(y_j w y_k^*)\|_2 < \varepsilon, \forall i \in \{1, 2\}, \forall j, k \in \{1, \dots, n\}$.

For the next part of the proof we need to introduce some notation. Thus, for each $i \in \{1, 2\}$ let $\{\xi_i^j\}_j \subset M_i$ be a BHOB of M_i over B . Also take $\{\xi_3^j\}_j = \{u_g\}_{g \in \mathbb{F}_2} \subset L(\mathbb{F}_2)$

such that $\xi_3^0 = u_e = 1$. Then using the notation $\tilde{M} = M *_B (B \otimes L(\mathbb{F}_2))$ as in Proposition 3.3 a simple exercise shows that

$$\beta = \{1\} \cup \{\xi_{i_1}^{j_1} \cdots \xi_{i_n}^{j_n} \mid n \in \mathbb{N}, i_k \in \{1, 2, 3\}, j_k \geq 1, i_1 \neq i_2, \dots, i_{n-1} \neq i_n\}$$

is BHOB of $\tilde{M}$ over B .

For each $n_0 \in \mathbb{N}$ let

$$S_{n_0} = \{1\} \cup \{\xi_{i_1}^{j_1} \cdots \xi_{i_n}^{j_n} \mid n \leq n_0, i_k \in \{1, 2\}, 1 \leq j_k \leq n_0, i_1 \neq i_2, \dots, i_{n-1} \neq i_n\},$$

$$\tilde{S}_{n_0} = \{1\} \cup \{\xi_{i_1}^{j_1} \cdots \xi_{i_n}^{j_n} \mid n \leq n_0, i_k \in \{1, 2, 3\}, 1 \leq j_k \leq n_0, i_1 \neq i_2, \dots, i_{n-1} \neq i_n\}.$$

Also for $i \in \{1, 2\}$ let $\tilde{S}_{n_0}^{R,i}$ (resp. $\tilde{S}_{n_0}^{L,i}$) be the subset of $\tilde{S}_{n_0}$ consisting of 1 and the vectors in $\tilde{S}_{n_0}$ such that $i_n \neq i$ (resp. $i_1 \neq i$). Note that if $\zeta, \zeta' \in \tilde{S}_{n_0}^{R,i}$ and $x \in M_i$ such that $E_B(x) = 0$, then $\|\zeta' x \zeta^*\|_2 = \|x\|_2$. Also if $\zeta \in \tilde{S}_{n_0}$ and $b \in B$ then $\|\zeta b\|_2 = \|b\|_2$.

We now strengthen *Fact 1* so that the elements $y_1, \dots, y_n$ may be taken in $\tilde{M}$:

Fact 2: $\forall \varepsilon > 0, \forall y_1, \dots, y_n \in \tilde{M}, \exists w \in \mathcal{U}(Q)$ such that $\|E_{M_i}(y_j^* w y_k)\|_2 < \varepsilon, \forall i \in \{1, 2\}, \forall j, k \in \{1, \dots, n\}$.

As our basis for $\tilde{M}$ is made up of bounded vectors by first approximating the y_k 's on the right of w and then approximating the y_j 's on the left of w we may assume that all of the y_j 's are basis elements and then use the triangle inequality to deduce the general case. Also as E_{M_i} is B -bimodular it enough to suppose that the y_j 's all lie in β . Thus we only need to show that $\forall \varepsilon > 0, \forall n_0 \in \mathbb{N}, \exists w \in \mathcal{U}(Q)$ such that $\|E_{M_i}(\zeta^* w \zeta')\|_2 < \varepsilon, \forall \zeta, \zeta' \in \tilde{S}_{n_0}$.

To prove this, we first use *Fact 1* to deduce that there exists $w \in \mathcal{U}(Q)$ such that $\|E_{M_i}(\zeta_0^* w \zeta'_0)\|_2 < \varepsilon, \forall \zeta_0, \zeta'_0 \in S_{n_0}$. Then if $\zeta, \zeta' \in \tilde{S}_{n_0}$ we may find $\zeta_1, \zeta'_1 \in \tilde{S}_{n_0}^{L,1} \cap \tilde{S}_{n_0}^{L,2}$, and $\zeta_0, \zeta'_0 \in S_{n_0}$ such that $\zeta = \zeta_0 \zeta_1$, and $\zeta' = \zeta'_0 \zeta'_1$. If $\zeta_1 = \zeta'_1 = 1$ then from above we have $\|E_{M_i}(\zeta^* w \zeta')\|_2 = \|E_{M_i}(\zeta_0^* w \zeta'_0)\|_2 < \varepsilon$. Otherwise we have $\|E_{M_i}(\zeta^* w \zeta')\|_2 \leq \|E_M(\zeta^* w \zeta')\|_2 \leq \|\zeta_1^* E_B(\zeta_0^* w \zeta'_0) \zeta'_1\|_2 \leq \|E_{M_1}(\zeta_0^* w \zeta'_0)\|_2 < \varepsilon$. This proves *Fact 2*.

We continue by showing that there are elements of $\mathcal{U}(Q)$ ("at infinity") which are almost orthogonal to the subspaces having at most one "large letter" from M_1, M_2 . Specifically, let $\mathcal{H}_{n_0} = \overline{\text{sp}}(\tilde{S}_{n_0} M_1 \tilde{S}_{n_0}^* \cup \tilde{S}_{n_0} M_2 \tilde{S}_{n_0}^*) \subset L^2(\tilde{M})$.

Suppose $\zeta_1, \zeta'_1 \in \tilde{S}_{n_0}^{R,i}, \zeta_2, \zeta'_2 \in \tilde{S}_{n_0}^{R,j}$, then $\forall b_1, b_2 \in B, \forall K, L > n_0$ we have that $E_B((\zeta'_1 \xi_i^K b_1 \zeta_1^*)^* (\zeta'_2 \xi_j^L b_2 \zeta_2^*)) = \delta_{\zeta'_1 \zeta'_2} E_B((\xi_i^K b_1 \zeta_1^*)^* (\xi_j^L b_2 \zeta_2^*))$, and also we have that $E_B((\xi_j^L b_2 \zeta_2^*)(\xi_i^K b_1 \zeta_1^*)^*) = \delta_{\zeta_1 \zeta_2} E_B((\xi_j^L b_2)(\xi_i^K b_1)^*)$. Hence

$$\begin{aligned} \langle \zeta'_1 \xi_i^K b_1 \zeta_1^*, \zeta'_2 \xi_j^L b_2 \zeta_2^* \rangle &= \tau \circ E_B((\zeta'_1 \xi_i^K b_1 \zeta_1^*)^* (\zeta'_2 \xi_j^L b_2 \zeta_2^*)) \\ &= \delta_{\zeta'_1 \zeta'_2} \tau \circ E_B((\xi_i^K b_1 \zeta_1^*)^* (\xi_j^L b_2 \zeta_2^*)) = \delta_{\zeta'_1 \zeta'_2} \delta_{\zeta_1 \zeta_2} \delta_{ij} \langle \xi_i^K b_1, \xi_j^L b_2 \rangle. \end{aligned}$$

Also if $\zeta \in \tilde{S}_{n_0}$, and $b \in B$ then $\langle \zeta'_1 \xi_i^K b_1 \zeta_1^*, \zeta b \rangle = \tau \circ E_B((\zeta b)^* \zeta'_1 \xi_i^K b_1 \zeta_1^*) = 0$.

Hence we have the following direct sum of orthogonal subspaces of $\mathcal{H}_{n_0}$:

$$\mathcal{H}'_{n_0} = \bigoplus_{\zeta \in \tilde{S}_{n_0}} \mathcal{H}_\zeta \oplus \bigoplus_{i \in \{1,2\}, \zeta, \zeta' \in \tilde{S}_{n_0}^{R,i}} \mathcal{H}_{i, \zeta', \zeta},$$

where $\mathcal{H}_\zeta = \overline{\zeta B}$, and $\mathcal{H}_{i, \zeta', \zeta} = \overline{\text{sp}}\{\xi_i^K\}_{K > n_0} B \zeta^*$.

Since $\{\xi_3^j\}_j = \{u_g\}_{g \in \mathbb{F}_2}$ we may find $m_0 > 2n_0$ such that $\{\xi_3^j\}_{j \leq n_0} \{\xi_3^j\}_{j \leq n_0}^* \subset \{\xi_3^j\}_{j \leq m_0}$. We will show that $\mathcal{H}_{n_0} \subset \mathcal{H}'_{m_0}$.

Let $\mathcal{K}_0$ be the Hilbert space generated by all vectors of the form $\eta' b \eta^*$ where $b \in B$, $\eta = \xi_{i_1}^{j_1} \cdots \xi_{i_n}^{j_n}$, $\eta' = \xi_{k_1}^{l_1} \cdots \xi_{k_m}^{l_m}$, such that $i_n \neq k_m$, $n + m \leq m_0$, and $j_p, l_p \leq m_0$, $\forall p$. If $\zeta, \zeta' \in \tilde{S}_{n_0}$, and $x \in M_i \ominus B$, Then we may find $\zeta_1, \zeta'_1 \in \tilde{S}_{n_0}^{R,i}$, and $\zeta_0, \zeta'_0 \in \tilde{S}_{n_0} \cap M_i$ such that $\zeta = \zeta_1 \zeta_0$, and $\zeta' = \zeta'_1 \zeta'_0$. We have that if P is the projection onto the subspace $\overline{\text{sp}}\{\xi_i^j\}_{j=1}^{m_0} B$ then $\zeta'_1(\zeta'_0 x \zeta_0^* - P(\zeta'_0 x \zeta_0^*) - E_B(\zeta'_0 x \zeta_0^*)) \zeta_1^* \in \mathcal{H}'_{m_0}$, $\zeta'_1 P(\zeta'_0 x \zeta_0^*) \zeta_1^* \in \mathcal{K}_0$, and $\zeta'_1 E_B(\zeta'_0 x \zeta_0^*) \zeta_0^* \in \tilde{S}_{n_0} B \tilde{S}_{n_0}^*$.

If $\zeta, \zeta' \in \tilde{S}_{n_0}$ and $b \in B$ then if ζ and ζ' do not end with a letter in the same algebra we have that $\zeta' b \zeta^* \in \mathcal{K}_0$, also if both ζ and ζ' end with something in $\{\xi_3^j\}_j$ then as $L(\mathbb{F}_2)$ commutes with B and since $\{\xi_3^j\}_{j \leq n_0} \{\xi_3^j\}_{j \leq n_0}^* \subset \{\xi_3^j\}_{j \leq m_0}$ we may rewrite $\zeta' b \zeta^*$ to see that it is in $\mathcal{K}_0$, otherwise as above we may find $\zeta_1, \zeta'_1 \in \tilde{S}_{n_0}^{R,i}$, and $\zeta_0, \zeta'_0 \in \tilde{S}_{n_0} \cap M_i$ such that $\zeta = \zeta_1 \zeta_0$, and $\zeta' = \zeta'_1 \zeta'_0$, and then decompose $\zeta' b \zeta^*$ into parts in $\mathcal{H}'_{m_0}$, $\mathcal{K}_0$, and something in $\tilde{S}_{n_0} B \tilde{S}_{n_0}$ with shorter words. Hence by induction to show that $\mathcal{H}_{n_0} \subset \mathcal{H}'_{m_0}$ it is enough to show that $\mathcal{K}_0 \subset \mathcal{H}'_{m_0}$.

Let η and η' be as above, and take $b \in B$, if $n = 0$, i.e. $\eta = 1$ then $\eta' b \eta^* = \eta' b \in \mathcal{H}_{m_0}$. Also if $i_n = 3$ then since $L(\mathbb{F}_2)$ commutes with B and $\xi_{i_n}^{j_n} \in \{\xi_3^j\}_{j \leq m_0}$ we can rewrite $\eta' b \eta^*$ so that η and η' are still in $\tilde{S}_{m_0}$ but such that the length of η is shorter. If $i_n = i \in \{1, 2\}$ then since $E_B(b \xi_{i_n}^{j_n}) = 0$ as above we may replace $b \xi_{i_n}^{j_n}$ with $P(b \xi_{i_n}^{j_n})$ and $b \xi_{i_n}^{j_n} - P(b \xi_{i_n}^{j_n})$ and in so doing rewrite $\eta' b \eta^*$ as a sum of things where the word on the right has shorter length plus something in $\mathcal{H}'_{m_0}$. Thus by induction we have shown that $\mathcal{K}_0 \subset \mathcal{H}'_{m_0}$ and so $\mathcal{H}_{n_0} \subset \mathcal{H}'_{m_0}$.

Let P_{n_0} be the orthogonal projection of $L^2(\tilde{M})$ onto $\mathcal{H}_{n_0}$.

Fact 3: $\forall \varepsilon > 0$, $y_1, \dots, y_n \in \tilde{M}$, $\exists w \in \mathcal{U}(Q)$ such that $\|P_{n_0}(y_j^* w y_k)\|_2^2 < \varepsilon$, $\forall j, k$.

Let m_0 be as above, then by *Fact 2* $\exists w \in \mathcal{U}(Q)$ such that $\|E_{M_i}(\zeta^* y_k^* w^* y_j \zeta')\|_2^2 < \varepsilon/3 |\tilde{S}_{m_0}|^2$, $\forall j, k \leq n$, $\forall \zeta, \zeta' \in \tilde{S}_{m_0}$.

Thus $\forall \zeta \in \tilde{S}_{m_0}$, $b \in B$ we have

$$\begin{aligned} |\langle \zeta b, y_j^* w y_k \rangle|^2 &= |\tau(E_B(y_k^* w^* y_j \zeta) b)|^2 \\ &\leq \|E_{M_1}(y_k^* w^* y_j \zeta)\|_2^2 \|b\|_2^2 < (\varepsilon/3 |\tilde{S}_{m_0}|^2) \|\zeta b\|_2^2, \end{aligned}$$

and so $\|P_{\mathcal{H}_\zeta}(y_j^* w y_k)\|_2^2 < \varepsilon/3 |\tilde{S}_{m_0}|^2$. Also $\forall i \in \{1, 2\}$, $\zeta, \zeta' \in \tilde{S}_{m_0}^{R,i}$, and $\xi \in \overline{\text{sp}}\{\xi_i^K\}_{K > m_0} B$ we have

$$|\langle \zeta' \xi \zeta^*, y_j^* w y_k \rangle|^2 = |\tau(E_{M_i}(\zeta^* y_k^* w^* y_j \zeta') \xi)|^2$$

$$\leq \|E_{M_i}(\zeta^* y_k^* w^* y_j \zeta')\|_2^2 \|\xi\|_2^2 < (\varepsilon/3|\tilde{S}_{m_0}|^2) \|\zeta' \xi \zeta^*\|_2^2,$$

and so $\|P_{\mathcal{H}_{i,\zeta',\zeta}}(y_j^* w y_k)\|_2^2 < \varepsilon/3|\tilde{S}_{m_0}|^2$.

Therefore

$$\begin{aligned} \|P_{n_0}(y_j^* w y_k)\|_2^2 &\leq \|P_{\mathcal{H}'_{m_0}}(y_j^* w y_k)\|_2^2 \\ &= (\sum_{\zeta \in \tilde{S}_{m_0}} \|P_{\mathcal{H}_\zeta}(y_j^* w y_k)\|_2^2) + (\sum_{i \in \{1,2\}, \zeta' \in \tilde{S}_{m_0}^{R,i}, \zeta \in \tilde{S}_{m_0}^{L,i}} \|P_{\mathcal{H}_{i,\zeta',\zeta}}(y_j^* w y_k)\|_2^2) < \varepsilon, \end{aligned}$$

for each $j, k \leq n$, thus we have proved *Fact 3*.

Next we note that if $Q' \cap \langle M, e_B \rangle$ contains a non-zero, finite-trace projection then so does $Q' \cap \langle M, e_{M_1} \rangle$ and so by our assumption we are in the position of applying Proposition 3.3.

Hence there exists a non-zero partial isometry $v \in \tilde{M}$ such that $vy = \theta(y)v$, $\forall y \in Q$. Let $\varepsilon > 0$ and take n_1 large enough so that there exists $v_0^* \in \text{sp} \tilde{S}_{n_1} B$ satisfying $\|v - v_0\|_2 < \varepsilon/6$. Using the notation in Proposition 3.4 let $L \in \mathbb{N}$, such that $\|\hat{E}_L(x)\|_2 < \varepsilon/12\|v_0\|_2^2$, $\forall x \in Q$, $\|x\| \leq 1$, also let $n_0 = n_1 + 3L$, and let m_0 be as above so that $\mathcal{H}_{n_0} \subset \mathcal{H}'_{m_0}$. Then as our basis is bounded we have that v_0 is bounded so by *Fact 3* $\exists w \in \mathcal{U}(Q)$, such that $\|P_{m_0}(v_0 w)\|_2 < \varepsilon/4\|v_0\|$. Take $w_0 \in M$ such that $\hat{E}_L(w_0) = 0$, and $\|w - w_0\|_2 < \varepsilon/6\|v_0\|_2^2$.

Let $\mathcal{K} \subset L^2(\tilde{M})$ be the right Hilbert B -module generated by 1 and all the vectors $\xi_{i_1}^{j_1} \dots \xi_{i_n}^{j_n} \in \beta$ such that if $i_{k_0} = 3$ for some $k_0 \leq n$ then $j_k \leq n_1$, $\forall k < k_0$. Note that $\mathcal{KM} \subset \mathcal{K}$ and so since $w_0 \in M$ and $v_0 \in \mathcal{K}$ we have that $v_0 w_0 \in \mathcal{K}$.

Let $\zeta, \zeta' \in \tilde{S}_{m_0}^{R,i}$, $K > m_0$, and $b \in B$, then since $K > n_1$ we have that $P_{\mathcal{K}}(\zeta' \xi_i^K b \zeta) = \zeta' \xi_i^K b \zeta$ if $\zeta \in M$ and $\zeta' \in \mathcal{K}$, and $P_{\mathcal{K}}(\zeta' \xi_i^K b \zeta) = 0$ otherwise. Also if $\zeta \in \tilde{S}_{m_0}$ and $B \in B$ then $P_{\mathcal{K}}(\zeta b) = \zeta b$ if $\zeta \in \mathcal{K}$ and $P_{\mathcal{K}}(\zeta b) = 0$ otherwise. Hence $P_{\mathcal{K}}(\mathcal{H}_{n_0}) \subset P_{\mathcal{K}}(\mathcal{H}'_{m_0}) \subset \mathcal{H}'_{m_0} \subset \mathcal{H}_{m_0}$.

Let us write v_0^* and w_0 in β as $v_0^* = \sum_{\xi_v \in \beta} \xi_v b_{\xi_v}$ and $w_0 = \sum_{\xi_w \in \beta} \xi_w b_{\xi_w}$. Take $\xi_v, \xi_w \in \beta$ such that $b_{\xi_v} \neq 0$ and $b_{\xi_w} \neq 0$, where $\xi_v = \xi_{i_1}^{j_1} \dots \xi_{i_n}^{j_n}$, $\xi_w = \xi_{k_1}^{l_1} \dots \xi_{k_m}^{l_m}$. Thus $\theta(\xi_w b_{\xi_w})(\xi_v b_{\xi_v})^* = (u_{k_1} \xi_{k_1}^{l_1} u_{k_1}^* \dots u_{k_m} \xi_{k_m}^{l_m} u_{k_m}^* b_{\xi_w})(b_{\xi_v}^* \xi_{i_n}^{j_n} \dots \xi_{i_1}^{j_1})$. Let us assume that $n \geq 3m$ by adding on 1's at the end of this word if necessary. If $k_s \leq n_0$, $\forall 1 \leq s < m$ then since $v_0^* \in \text{sp} \tilde{S}_{n_1} B$, and $m < L$, by decomposing $u_{k_m}^* b_{\xi_w} b_{\xi_v}^* \xi_{i_n}^{j_n}$ as it's expectation onto B plus something with terms in $B \otimes L(\mathbb{F}_2)$ and zero expectation onto B , we write $\theta(\xi_w b_{\xi_w})(\xi_v b_{\xi_v})^*$ as something in $\mathcal{H}_{n_0}$ plus something in $\mathcal{K}^\perp$. Otherwise if $k_s > n_0$ for some $s < m$ then by decomposing $u_{k_m}(\xi_{k_m}^{l_m}(u_{k_m}^* b_{\xi_w} b_{\xi_v}^* \xi_{i_n}^{j_n}) \xi_{i_{n-1}}^{j_{n-1}}) \xi_{i_{n-2}}^{j_{n-2}}$ just as above into it's expectation onto B plus something with terms in $B \otimes L(\mathbb{F}_2)$ and zero expectation onto B , we write $\theta(\xi_w b_{\xi_w})(\xi_v b_{\xi_v})^*$ as something with shorter words plus something in $\mathcal{K}^\perp$. Hence by induction we have shown that $\theta(\xi_w b_{\xi_w})(\xi_v b_{\xi_v})^* \in \mathcal{H}_{n_0} + \mathcal{K}^\perp$ and hence also $\theta(w_0)v_0 \in \mathcal{H}_{n_0} + \mathcal{K}^\perp$.

As $P_{\mathcal{K}}(\mathcal{H}_{n_0}) \subset \mathcal{H}_{m_0}$ we have that $P_{\mathcal{K}}(\theta(w_0)v_0) \subset \mathcal{H}_{m_0}$. Thus

$$\begin{aligned} |\langle v_0 w_0, \theta(w_0)v_0 \rangle| &= |\langle v_0 w_0, P_{\mathcal{K}}(\theta(w_0)v_0) \rangle| = |\langle P_{m_0}(v_0 w_0), P_{\mathcal{K}}(\theta(w_0)v_0) \rangle| \\ &\leq \|P_{m_0}(v_0 w_0)\|_2 (\|w_0 - w\|_2 \|v_0\| + \|v_0\|_2) \\ &\leq (\|P_{m_0}(v_0 w)\|_2 + \|v_0\| \|w_0 - w\|_2) (\|w_0 - w\|_2 \|v_0\| + \|v_0\|_2) \end{aligned}$$

$$< (\varepsilon/4\|v_0\| + \varepsilon/6\|v_0\|)(\varepsilon/6\|v_0\| + \|v_0\|) < \varepsilon/4.$$

Hence we have shown that

$$\begin{aligned} \|v\|_2^2 &= \|vw\|_2^2 \\ &= \langle vw, \theta(w)v \rangle \leq 2\|v - v_0\|_2 + |\langle v_0w, \theta(w)v_0 \rangle| \\ &\leq 2\|v - v_0\|_2 + 2\|v_0\|^2\|w - w_0\|_2 + |\langle v_0w_0, \theta(w_0)v_0 \rangle| \\ &< \varepsilon/3 + \varepsilon/3 + \varepsilon/4 < \varepsilon, \end{aligned}$$

which contradicts the assumption that v is non-zero. $\square$

5. RIGID SUBALGEBRAS IN AFP FACTORS: GENERAL BASS-SERRE TYPE RESULTS

We have shown in Theorem 4.3 that if Q is a relatively rigid von Neumann subalgebra of an AFP algebra $M = M_1 *_B M_2$ then there exists a non-trivial Hilbert bimodule $\mathcal{H} \subset L^2(M)$ intertwining Q into one of the M_i 's. We now deduce that a corner of Q can be conjugated by a unitary element into that same M_i . When M_1, M_2 are factors, one can in fact uniquely partition 1 with projections $q_1, q_2 \in Q' \cap M$ such that Qq_i is unitary conjugate into M_i , $i = 1, 2$. This general Bass-Serre type result will be used in the next sections to derive more specific statements in the cases $B = \mathbb{C}$, $B = A$ abelian Cartan and $B = R$ the hyperfinite II_1 factor.

5.1. Theorem. *Let (M_i, τ_i) , $i = 0, 1, 2$, be finite von Neumann algebras with a common von Neumann subalgebra $B \subset M_i$, $i = 0, 1, 2$, such that $\tau_0|_B = \tau_1|_B = \tau_2|_B$, and such that the inclusions $B \subset M_i$ are homogeneous. Denote $M = M_0 *_B M_1 *_B M_2$. Let $Q \subset M$ be a relatively rigid diffuse von Neumann subalgebra. Assume no corner of Q can be embedded into M_0 inside M .*

1°. *There exist $i \in \{1, 2\}$, projections $q \in Q$, $q'' \in Q' \cap M$ with $qq'' \neq 0$ and a unitary element $u \in \mathcal{U}(M)$ such that $uqQqq''u^* \subset M_i$.*

2°. *If M_1, M_2 are factors then there exists a unique pair of projections $q_1, q_2 \in Q' \cap M$ such that $q_1 + q_2 = 1$ and $u_i(Qq_i)u_i^* \subset M_i$ for some unitaries $u_i \in \mathcal{U}(M)$, $i = 1, 2$. Moreover, these projections lie in the center of $Q' \cap M$.*

Proof. 1°. By Theorem 4.3 and (2.1 in [P6]) there exist $i \in \{0, 1, 2\}$, $q \in \mathcal{P}(Q)$, $p \in \mathcal{P}(M_i)$ non-zero projections, an isomorphism ψ of qQq into $pM_i p$ and a non-zero partial isometry $v \in M$ such that $vv^* \in (qQq)' \cap qMq$, $v^*v \in \psi(qQq)' \cap pMp$ and $xv = v\psi(x)$, $\forall x \in qQq$. By hypothesis, i cannot be equal to 0, thus $i \in \{1, 2\}$. Note that by shrinking q if necessary, we may assume $xv = 0$ for $x \in qQq$ implies $x = 0$. Also, if we denote by q' the support projection of $E_{M_1}(v^*v)$, then by replacing if necessary ψ by $q'\psi(\cdot)q'$ it follows that we may assume $q' = p$.

Now note that if a corner of $\psi(qQq)$ can be embedded into pBp inside $pM_i p$ then by Lemma 1.4.5 a corner of Q can be embedded into B (thus into $M_0 \supset B$ as well) inside M , contradicting the hypothesis. Thus, no corner of $\psi(qQq) \subset pM_i p$ can be embedded into B inside M_i , so we can apply Theorem 1.2.1 to conclude that $Q'_0 \cap pMp \subset pM_i p$. Hence,

$v^*v \in Q'_0 \cap pMp \subset M_i$. Taking $q'' = vv^*$ and $u \in M$ a unitary element such that $uqq'' = v$, the statement follows.

2°. Let $z = z(q)$ denote the central support of q in Q and note that zq'' is then the central support of qq'' in Qq'' . By the factoriality of M_i , $i = 1, 2$, it follows that there exists a unitary $u \in \mathcal{U}(M)$ such that $Qq''z \subset uM_iu^*$. (Indeed, this is because whenever $Q_0 \subset N$ is an inclusion of finite von Neumann algebras, $q_0 \in \mathcal{P}(Q_0)$ and $N_0 \subset N$ is a subfactor with $qQ_0q \subset N_0$, then there exists $u \in \mathcal{U}(N)$ such that $Q_0z(q) \subset uN_0u^*$.) Thus, the projection $p'_0 = q''z \in Q' \cap M$ together with the unitary element u satisfy the condition $uQp'_0u^* \subset M_i$.

Let $\mathcal{F}$ be the set of all families of mutually orthogonal projections $\{p'_i\}_{i \in I} \subset \mathcal{P}(Q' \cap M)$, with the property that $\forall i \in I, \exists j(i) \in \{1, 2\}$ (unique by Theorem 1.2.1) and $v_i \in \mathcal{U}(M)$ such that $v_iQp'_iv_i^* \subset M_{j(i)}$. $\mathcal{F}$ is clearly inductively ordered with respect to the order given by inclusion. Let $\{p'_i\}_{i \in I}$ be a maximal element. Let $q'_1 = \sum_{j(i)=1} p'_i$, $q'_2 = \sum_{j(i)=2} p'_i$ and set $q' = 1 - q'_1 - q'_2$.

Assume $q' \neq 0$. Since Q is diffuse, there exists $q \in \mathcal{P}(Q)$ such that $\tau(q'q) = 1/n$ for some integer $n \geq 1$. Let $\tilde{Q} \subset M$ be a von Neumann algebra isomorphic to $M_{n \times n}(qQq')$ with qQq' equal to the upper-left corner $qq'\tilde{Q}qq'$ and qq' having central trace $1/n$ in $\tilde{Q}$. By (Sec. 4 in [P5]) $\tilde{Q} \subset M$ is a rigid inclusion. Thus, we can apply the first part of the proof to get $i \in \{1, 2\}$, $0 \neq \tilde{q}' \in \tilde{Q}' \cap M$ and a unitary element $w \in M$ such that $w\tilde{Q}\tilde{q}'w^* \subset M_i$. Since qq' has scalar central trace in $\tilde{Q}$, it follows that the projection $p = qq'\tilde{q}' \in (qQq')' \cap qMq = q(Q' \cap M)q$ is non-zero. Thus $p = qq''$, for some projection $q'' \in Q' \cap M$ with $q'' \leq q'$. Since $p \leq \tilde{q}'$, we also have $w(qQq'')w^* \subset M_i$, implying that if we denote $p' = z(q)q''$ (where $z(q)$ is the central support of q in Q) then $p' \in Q' \cap M$, $p' \leq q'$ and there exists a unitary element u in M such that $u(Qp')u^* \subset M_i$. But then $\{p'_i\}_{i \in I} \cup \{p'\}$ lies in $\mathcal{F}$, thus contradicting the maximality of $\{p'_i\}_{i \in I}$.

We have thus shown that $q'_1 + q'_2 = 1$. On the other hand, by the factoriality of the M_k 's, $k = 1, 2$, for each fixed k we can choose the unitary elements $\{v_i \mid j(i) = k\}$ which satisfy $v_i(Qp'_i)v_i^* \subset M_k$ so that $v_ip'_iv_i^*$ be mutually orthogonal projections in M_k . Taking $u_k \in \mathcal{U}(M)$ to be a unitary element extending $\sum_{j(i)=k} v_i$, it follows that $u_k(Qq'_k)u_k^* \subset M_k$, $k = 1, 2$.

This proves the existence part of part 2°. But the uniqueness part is then clear, since if p'_1, p'_2 is another pair of projections in $Q' \cap M$ satisfying $p'_1 + p'_2 = 1$, $v_i(Qp'_i)v_i^* \subset M_i$ for some $v_i \in \mathcal{U}(M)$, $i = 1, 2$, and we assume $x = p'_1q'_2 \neq 0$, then the partial isometry w in the polar decomposition of x lies in $Q' \cap M$ and if we denote $p = ww^*$ then $v_1(Qp)v_1^* \subset M_1$ while $u_2w^*(Qp)wu_2^* \subset M_2$, contradicting Theorem 1.1.

To finish the proof of 2° we need to show that q'_1, q'_2 are in the center of $Q' \cap M$. Since $q'_1 + q'_2 = 1$, this amounts to showing that their central supports in $Q' \cap M$ are disjoint.

Assume by contradiction that there exist non-zero projections $q''_i \leq q'_i$, $q''_i \in Q' \cap M$ with $u'q''_iu'^* = q''_i$ for some $u' \in \mathcal{U}(Q' \cap M)$. But then $u_k(Qq''_k)u_k^* \subset M_k$, $k = 1, 2$ are diffuse and are conjugate by the unitary element $u_2u'u_1^*$, contradicting Theorem 1.1 again. $\square$

5.2. Theorem. *Let I be a set of indices with $0 \in I$ and (M_i, τ_i) , $i \in I$, a family of finite von Neumann algebras with a common von Neumann subalgebra $B \subset M_i$, such that $\tau_{0|B} =$*

$\tau_i|_B, \forall i$. Assume M_i are factors for $i \neq 0$ and the inclusions $B \subset M_i$ are homogeneous, $\forall i \in I$. Denote $M = *_{B, i \in I} M_i$ the free product with amalgamation over B of the algebras $M_i, i \in I$. Let $t > 0$ and $Q \subset M^t$ be a relatively rigid diffuse von Neumann subalgebra such that no corner of Q can be embedded into M_0 inside M and such that the normalizer of Q in M^t generates a factor N . Then there exists a unique $i \in I \setminus \{0\}$ and a unitary element $u \in M^t$ such that $uQu^* \subset M_i^t$. Moreover, such u satisfies $uNu^* \subset M_i^t$, and in fact $u\tilde{N}u^* \subset M_i^t$, where $\tilde{N} = \tilde{N}(N, M_1^t; B)$ is as defined in 1.2.2.

Proof. Note first that $Q \subset M^t$ rigid implies Q is countably generated (see e.g. [PeP]). Thus, there exists a countable subset $S \ni 0$ of indices $i \in I$ such that $Q \subset (*_{B, i \in S} M_i)^t$. By 3.2, $Q \subset (*_{B, i \in S} M_i)^t$ is rigid and by 1.2.1 all N is contained in $(*_{B, i \in S} M_i)^t$. This shows that it is sufficient to prove the statement in the case $M_i, i \geq 0$, is a sequence of algebras.

Moreover, since $Q \subset M^t$ is rigid and since the factors $\tilde{M}(K, t) = (M_0 *_B (*_{B, k \in K} M_k))^t$, with K finite subset of $\{1, 2, \dots\}$, tend to M^t , it follows by ([P5]) that there exists a non-zero projection $q' \in Q' \cap M^t$, a unitary element $v \in M^t$ and a finite set $K \subset \{1, 2, \dots\}$ such that $v(Qq')v^* \subset \tilde{M}(K, t)$. But $q' \in N$ and by (3.4 in [P6]) Qq' is quasi-regular in $q'Nq'$, so by 1.2.1 we have $v(q'Nq')v^* \subset \tilde{M}(K, t)$. Since N is a factor, v can be modified so that $vNv^* \subset \tilde{M}(K, t)$. In particular $vQv^* \subset \tilde{M}(K, t)$.

Since Q is diffuse, there exists $q \in \mathcal{P}(Q)$ such that $\tau(q) \leq t^{-1}$. Thus, we may assume $v(qQq)v^* \subset \tilde{M}_K \stackrel{\text{def}}{=} M_0 *_B (*_{B, k \in K} M_k)$, and notice that by (4.7 in [P5]) the unital inclusion $v(qQq)v^* \subset p\tilde{M}_K p$ is rigid, where $p = vqv^*$. Since K is finite, 5.1 applies to get $i \in K$, $0 \neq q'_i \in (vqQqv^*)' \cap p\tilde{M}_K p$ and a unitary $w \in \tilde{M}_K$ such that $w(vqQqv^*q'_i)w^* \subset M_i$. Moreover, since $\tau(q) \leq t^{-1}$ we can view $w(vqQqv^*q'_i)w^*$ as a (possibly non-unital) subalgebra of M_i^t . Since $q'_i \in (vqQqv^*)' \cap pMp$, it follows that $q'_i \in vqNqv^*$ (recall that N is generated by the normalizer of Q in M). By (3.4 in [P6]) and 1.2.1 again, it follows that $w(q'_i vqNqv^* q'_i)w^* \subset M_i$, implying that $wq'_i vq$ can be extended to a unitary element $u \in M^t$ such that $uNu^* \subset M_i^t$, thus $uQu^* \subset M_i^t$. Also, by 1.2.1 i is unique with this property while by Corollary 1.2.3 it follows that $u\tilde{N}u^* \subset M_i^t$. $\square$

6. AMALGAMATION OVER $\mathbb{C}$: FREE PRODUCT FACTORS WITH PRESCRIBED $\mathcal{F}(M)$.

We first apply Theorem 5.1 to plain free product factors, where the result becomes an analogue of the classical Kurosh theorem for groups. The first Kurosh-type results in Operator Algebra framework were obtained by N. Ozawa in ([O2]). He proved that if N is a non-prime non-hyperfinite II_1 subfactor of a free product $M = M_1 * M_2$ of semiexact finite factors M_1, M_2 , then N can be unitarily conjugated into either M_1 or M_2 (this is analogue to “Kurosh subgroup theorem”). As a consequence, he showed that if two free products $*_i M_i, *_j N_j$ of non-hyperfinite, non-prime, semiexact factors N_i, M_j are isomorphic then the “length” of the two free products must be the same and each N_i is unitary conjugate to M_i , after some permutation of indices (this is analogue to “Kurosh isomorphism theorem”).

In turn, our results cover different classes of algebras. Thus, our analogue of the “Kurosh subgroup theorem” allows M_1, M_2 to be arbitrary finite von Neumann algebras but only

gives information about relatively rigid subalgebras Q of $M_1 * M_2$. Our corresponding “isomorphism theorem”, which in fact we obtain for amplifications of free products, will require the factors N_i, M_j to be either w-rigid, i.e. to have diffuse, regular, relatively rigid subalgebras, or to be group measure space factors associated to actions of w-rigid ICC groups. In particular, it holds true for N_i, M_j II_1 factors with the property (T) (in the sense of [CJ]) and more generally for tensor products of property (T) II_1 factors with arbitrary finite factors. Moreover, since the factors $N = L(\mathbb{Z}^2 \rtimes \mathbb{F}_n)$ in ([P5]) are w-rigid and have trivial fundamental group (see Theorem 7.11 for a different proof), this will allow us to obtain large classes of factors with trivial fundamental group, different from the ones in ([P5,6]). More generally, using also ([DyR]), we construct a completely new class of factors with prescribed fundamental group which, unlike the ones in ([P5,6]), have no Cartan subalgebras (by [V2], cf. 6.6 below).

6.1. Theorem. *Let (M_i, τ_i) , $i = 0, 1, 2$ be finite von Neumann algebras and denote $M = M_0 * M_1 * M_2$. Assume no direct summand of (M_0, τ_0) has relatively rigid diffuse von Neumann subalgebras (e.g. $M_0 = \mathbb{C}$, or more generally M_0 with Haagerup property). Let $Q \subset M$ be a relatively rigid diffuse von Neumann subalgebra of M .*

1°. *There exist $i \in \{1, 2\}$, $q \in \mathcal{P}(Q)$, $q' \in \mathcal{P}(Q' \cap M)$ and $u \in \mathcal{U}(M)$ such that $qq' \neq 0$ and $uqQqq'u^* \subset M_i$.*

2°. *If in addition M_1, M_2 are factors, then there exists a unique pair of projections $q'_1, q'_2 \in Q' \cap M$ such that $q'_1 + q'_2 = 1$ and $u_i(Qq'_i)u_i^* \subset M_i$ for some unitaries $u_i \in \mathcal{U}(M)$, $i = 1, 2$. Moreover, $q'_1, q'_2 \in \mathcal{Z}(Q' \cap M)$.*

3°. *If instead of M_1, M_2 we consider a whole family of finite factors $M_i, i \geq 1$, and we take $Q \subset M^t = (M_0 * M_1 * M_2 * \dots)^t$ a rigid inclusion, for some $t > 0$, and we assume the normalizer of Q in M^t generates a factor N , then there exists a unique $i \geq 1$ and a unitary element $u \in M^t$ such that $uQu^* \subset M_i^t$. Moreover, such u satisfies $uNu^* \subset M_i^t$, and in fact $u\tilde{N}u^* \subset M_i^t$, where $\tilde{N} = \tilde{N}(N, M_i^t; \mathbb{C})$ is as defined in 1.2.2.*

Proof. As in the proof of 5.2, note that Q relatively rigid implies Q countably generated. Thus, there exist countably generated von Neumann subalgebras $M_i^0 \subset M_i, i = 0, 1, 2$, such that $Q \subset M_0^0 * M_1^0 * M_2^0$. Thus, to prove 1° it is clearly sufficient to prove it in the case M_i are countable generated, $i = 0, 1, 2$. But then each $\mathbb{C} \subset M_i$ is homogeneous by Lemma 4.2. Let us show that no corner of Q can be embedded into M_0 inside M . Assume this is not true. By (2.1. in [Po6]) it follows that there exist $q \in Q, p \in M_0$ non-zero projections, a unital isomorphism ψ of qQq into pM_0p and a non-zero partial isometry $v \in M$ such that $vv^* \in (qQq)' \cap qMq, v^*v \in \psi(qQq)' \cap pMp$ and $xv = v\psi(x), \forall x \in qQq$. Denote $q'' = vv^* \in Q' \cap M$.

Since $\psi(qQq) \subset pM_0p$ is a diffuse von Neumann subalgebra, by Theorem 1.2.1, it follows that $\psi(qQq)' \cap pMp \subset pM_0p$. Thus $v^*v \in pM_0p$. This shows that $v^*qQqv \subset pM_0p$, which in turn implies that $qQqq'' \subset wM_0w^*$, for some unitary element $w \in M$ extending v . Since $Q \subset M$ is rigid, by (Proposition 4.7 in [Po5]) $qQqq'' \subset qq''Mqq''$ is also rigid, which trivially implies $qQqq'' \oplus (1 - qq'')\mathbb{C} \subset M$ is rigid. But then $w^*(qQqq'' \oplus (1 - qq'')\mathbb{C})w \subset M_0$ follows rigid by Corollary 3.2. By taking a suitable amplification of $w^*(qQqq'')w$ in M_0 and using

again (4.7 in [Po5]), this implies a direct summand of M_0 contains a relatively rigid diffuse von Neumann subalgebra, a contradiction.

Altogether, this shows that the conditions required in 5.1 are satisfied, so part 1° of the statement follows as a particular case of that theorem.

For part 2°, simply notice that if M_1, M_2 are factors then the countably generated von Neumann subalgebras M_0^0, M_1^0, M_2^0 with the property that $Q \subset M_0^0 * M_1^0 * M_2^0$ can be chosen so that M_1^0, M_2^0 are factors as well, so 5.1.2° applies.

Part 3° follows then from part 2° and 5.2. $\square$

6.2. Definition. A finite von Neumann algebra (M, τ) is *weakly rigid* (w-rigid) if it contains a regular, relatively rigid, diffuse von Neumann subalgebra, i.e. a subalgebra $Q \subset M$ such that $\mathcal{N}_M(Q)'' = M$ and $Q \subset M$ is a rigid inclusion (or Q is a relatively rigid subalgebra of M [P5]). Note that if G is a w-rigid group as defined in ([P5-8]), i.e. G contains an infinite normal subgroup with the relative property (T) of Kazhdan-Margulis, then $L(G)$ is w-rigid. Also, if M is w-rigid and P is an arbitrary finite von Neumann algebra then $M \overline{\otimes} P$ is w-rigid.

6.3. Theorem. *Let $(M_0, \tau_{M_0}), (N_0, \tau_{N_0})$ be finite von Neumann algebras which have no relatively rigid diffuse subalgebras (e.g. M_0, N_0 have Haagerup's compact approximation property). Let $M_1, \dots, M_m$ and $N_1, \dots, N_n$ be II_1 factors, where $n, m \geq 1$ are some cardinals (finite or infinite) and assume each M_i, N_j is w-rigid. If θ is an isomorphism of $M = *_{i=0}^m M_i$ onto N^t , where $N = *_{j=0}^n N_j$ and $t > 0$, then $m = n$ and, after some permutation of indices, $\theta(M_i)$ and N_i^t are unitary conjugate in N^t , $\forall i \geq 1$.*

Proof. For each $m \geq i \geq 1$ let $Q_i \subset M_i$ be a regular, relatively rigid, diffuse von Neumann subalgebra. Since M_i are factors and $Q_i \subset M_i$ are regular inclusions, it follows that for any $s > 0$ the factor M_i^s contains a regular, diffuse, relatively rigid subalgebra. To see this, note that $\mathcal{N}_{M_i}(Q_i)$ acts ergodically on the center of Q_i , so $\mathcal{Z}(Q_i)$ is either diffuse or atomic. In both cases we can find projections $q' \in \mathcal{Z}(Q_i)$ and $q \in Q_i q'$ such that the central trace of q in Q_i is a scalar multiple of q' and $\tau(q) = s/k$, for some $k \geq t$. By (Lemma 3.5 in [P6I]) it follows that the inclusion $qQ_i q \subset qM_i q$ is regular and by (4.7 on [P5]) it is rigid as well. But then $M_{k \times k}(qQ_i q) \subset M_{k \times k}(qM_i q) = M_i^s$ is regular and rigid ([P5]).

Moreover, note that any diffuse von Neumann subalgebra $B_i \subset M_i^s$ satisfies $B_i' \cap M^s \subset M_i^s$. To see this, note first that by taking direct sums of k copies of B_i embedded diagonally into $M_{k \times k}(M_i^s) = M_i^{ks}$, with k sufficiently large, we may assume $s \geq 1$. Then take $p \in \mathcal{P}(B_i)$ with $\tau(p) = 1/s$ and note that if we assume by contradiction that $B_i' \cap M^s \neq B_i' \cap M_i^s$, then $(pB_i p)' \cap pM^s p \neq (pB_i p)' \cap pM_i^s p$. But $(pM_i^s p \subset pMp) \simeq (M_i \subset M)$ and M splits off M_i as a free product. Thus, by Theorem 1.2.1 the relative commutant in $pM^s p = M$ of the diffuse subalgebra $pB_i p \subset pM_i^s p = M_i$ must be contained in $pM_i^s p$, contradicting the assumption.

Taking now $s = 1/t$, it follows that $M_i^{1/t}$ has a diffuse, regular, relatively rigid subalgebra, implying that $P_i = \theta^{1/t}(M_i^{1/t})$ has such a subalgebra B_i as well. In particular, the inclusion $B_i \subset N$ is rigid. In addition, $B_i' \cap N \subset P_i$. Since the inclusion $B_i \subset N$ is rigid and regular, by 6.1.3° there exists a unique $j(i) \in \{1, 2, \dots, n\}$ and a unitary $u_i \in N$ such that

$B_i \subset u_i N_{j(i)} u_i^*$ and $P_i \subset u_i N_{j(i)} u_i^*$. Thus, there exists a unique $j(i)$ such that for some unitary $v_i \in N^t$ we have $\theta(M_i) = P_i^t \subset v_i N_{j(i)}^t v_i^*$.

Similarly, by applying the above to θ^{-1} we get for each $n \geq j \geq 1$ a unique $m \geq k(j) \geq 1$ and a unitary element in $w_j \in M$ such that $\theta^{-1}(N_j^t) \subset w_j M_{k(j)} w_j^*$. Altogether, for each $m \geq i \geq 1$ we get

$$(6.3.1) \quad M_i = \theta^{-1}(\theta(M_i)) \subset \theta^{-1}(v_i N_{j(i)}^t v_i^*) \subset u_i M_{k(j(i))} u_i^*,$$

where $u_i = w_{j(i)} \theta^{-1}(v_i)$. By Theorem 1.2.1 it follows that $k(j(i)) = i$, i.e. $k \circ j = id$. Similarly, $j \circ k = id$. Thus $m = n$, j and k are onto isomorphisms and the inclusions (6.3.1) are in fact equalities. $\square$

In the next statement, for a finite permutation $\pi \in S_m$ and $m \geq i \geq 1$ we denote by $m(\pi, i)$ the cardinality of the set $\{\pi^k(i) \mid k \geq 1\}$.

6.4. Corollary. *Let $m \in \mathbb{N}$ and let $M_1, \dots, M_m$ be w -rigid II_1 factors. Let (M_0, τ_{M_0}) be a finite von Neuman algebra which contains no rigid diffuse von Neumann subalgebras. Denote $M = *_{i=0}^m M_i$. Then $\mathcal{F}(M) \subset \cup_{\pi \in S_m} \cap_{i=1}^m \mathcal{F}(M_i)^{m(\pi, i)^{-1}} \subset \cap_{i=1}^m \mathcal{F}(M_i)^{1/m!}$. In particular if one of the factors $M_i, m \geq i \geq 1$, has trivial fundamental group then so does M .*

Proof. For $t \in \mathcal{F}(M)$, let $\theta : M \rightarrow M^t$ be an isomorphism. Applying the previous theorem, we get that there exists $\pi \in S_m$ such that $\theta(M_i)$ and $M_{\pi(i)}^t$ are unitary conjugate in M^t . In particular, we have that $M_i \cong M_{\pi(i)}^t, \forall 1 \leq i \leq m$ which by induction implies that $M_i \cong M_{\pi^k(i)}^{t^k}, \forall 1 \leq i \leq m, k \in \mathbb{N}$.

Fixing i and letting $k = m(\pi, i)$ we obtain that $t^{m(\pi, i)} \in \mathcal{F}(M_i)$, or equivalently $t \in \mathcal{F}(M_i)^{m(\pi, i)^{-1}}$. By intersecting over all values of i , taking the union over all possible permutations and noticing that $m(\pi, i)$ divides m , the result follows. $\square$

For the next corollary we denote by $\mathcal{X}_{fin}$ the set of all finite tuples of positive numbers $(t_i)_{1 \leq i \leq n} \subset \mathbb{R}_+^*, n \geq 2$, and by $\mathcal{X}_\infty$ the set of all infinite sequences $(t_j)_{j \geq 1} \subset \mathbb{R}_+^*$. Also, we let $\mathcal{X} = \mathcal{X}_{fin} \cup \mathcal{X}_\infty$. If $X = (t_i)_i, Y = (s_j)_j$ are in $\mathcal{X}$ we then write $X \sim Y$ if both have the same “length” and there exists a permutation (bijection) π of the (common) set of indices $\{1, 2, \dots\}$ such that $s_{\pi(i)} = t_i, \forall i$.

Given a II_1 factor M and $X = (t_i)_i \in \mathcal{X}$, we denote $M^X = *_i M^{t_i}$. Note that if $X, Y \in \mathcal{X}$ and $X \sim_\pi Y$ then π implements a natural isomorphism $\theta_\pi : M^X \simeq M^Y$, in the obvious way. For $t > 0$ and $X = (t_i)_i \in \mathcal{X}$ we denote $tX = (tt_i)_i \in \mathcal{X}$.

6.5. Corollary. *Let M be a w -rigid II_1 factor with trivial fundamental group, e.g. $M = L(\mathbb{Z}^2 \rtimes SL(2, \mathbb{Z}))$ (cf. [P5]).*

1°. *If $X, Y \in \mathcal{X}$ then $M^X \simeq M^Y$ iff $X \sim Y$ and iff $M^X * L(\mathbb{F}_k) \simeq M^Y * L(\mathbb{F}_k)$, for some $1 \leq k \leq \infty$.*

2°. *$\mathcal{F}(M^X) = \{1\}$ for all $X \in \mathcal{X}_{fin}$. Moreover, if we denote by $\mathcal{X}_0$ the set of elements X in $\mathcal{X}_{fin}$ with $\min X = 1$, then $\{M^X \mid X \in \mathcal{X}_0\}$ is a continuous family of mutually non stably isomorphic II_1 factors.*

3°. For each $X \in \mathcal{X}_\infty$ denote $\mathcal{S}_X = \{t \in \mathbb{R}_+^* \mid tX \sim X\}$. Then $\mathcal{F}(M^X) = \mathcal{S}_X$. In particular, if $S \subset \mathbb{R}_+^*$ is an infinite, countable subgroup and $X \in \mathcal{X}_\infty$ has the elements of S as entries, each one repeated with the same (possibly infinite) multiplicity, then $\mathcal{F}(M^{tX}) = S$, $\forall t > 0$. Moreover, M^{t_1X}, M^{t_2X} are stably isomorphic iff t_1, t_2 are in the same class in $\mathbb{R}_+^*/S$. Thus, $\{M^{tX} \mid t \in \mathbb{R}_+^*/S\}$ is a continuous family of mutually non stably isomorphic II_1 factors all with fundamental group equal to S .

4°. If $S \subset \mathbb{R}_+^*$ is an arbitrary infinite (possibly uncountable) subgroup then the II_1 factor $M^S = *_{s \in S} M^s$ has fundamental group equal to S .

Proof. 1°. If $M^X \cong M^Y$ then by Theorem 6.2 X, Y have the same length and there exists a bijection π of the corresponding (finite or infinite) set of indices $\{1, 2, \dots\}$ such that $M^{t_i} = M^{s_{\pi(i)}}$, $\forall i$. Since M has trivial fundamental group, this implies $t_i = s_{\pi(i)}$, $\forall i$, and thus $X \sim_\pi Y$. The second equivalence has exactly the same proof, using 6.3 with $M_0 = L(\mathbb{F}_k)$.

2°. If $X = (t_i)_{1 \leq i \leq n}$, $Y = (s_j)_{1 \leq j \leq m}$ are in $\mathcal{X}_{fin}$ and $M^X \simeq (M^Y)^t$ then $M^X * L(\mathbb{F}_\infty) \simeq (M^Y)^t * L(\mathbb{F}_\infty)$. But by ([DyR]) the last factor is isomorphic to $M^{tY} * L(\mathbb{F}_\infty)$. Thus, $M^X * L(\mathbb{F}_\infty) \simeq M^{tY} * L(\mathbb{F}_\infty)$, which by part 1° implies $X \sim tY$. Thus, if $X = Y$ (respectively $X, Y \in \mathcal{X}_0$) then $t = 1$ and we get $\mathcal{F}(M^X) = \{1\}$ (resp. $X \sim Y$). This implies both statements.

3°. By ([DyR]), if $X \in \mathcal{X}_\infty$ then $(M^X)^t \cong M^{(tX)}$. Thus $M^X \cong (M^X)^t$ if and only if $X \sim tX$, which readily implies all statements.

4°. It is easy to see that the proof of the amplification formula $(M^S)^t \simeq M^{tS}$ in ([DyR]) doesn't in fact depend on the fact that the infinite set S is countable. Thus, since for $t \in S$ we have $tS = S$ as sets, it follows that $(M^S)^t = M^{tS} = M^S$, thus $S \subset \mathcal{F}(M^S)$. Conversely, if $s \in \mathbb{R}_+^*$ satisfies $(M^S)^s \simeq M^S$, then $M^{sS} \simeq M^S$, which by 6.3 implies $tS = S$, so that $s \in S$. $\square$

6.6. Remarks. Dima Shlyakhtenko pointed out to us that by combining Voiculescu's initial argument for showing that $L(\mathbb{F}_n)$ has no Cartan subalgebras, with Kenley Jung's "monotonicity" ([Ju]) it follows that any free product of type II_1 factors is "Cartan-less" (see [Sh] for a detailed argument). Thus, unlike the examples of factors with prescribed fundamental group in ([P6]), which are group measure space factors associated with equivalence relations coming from Connes-Størmer Bernoulli actions, the examples of factors M that we produce here (in Corollary 6.5) have no Cartan subalgebras, and altogether no diffuse hyperfinite "core". In particular, they cannot be written as crossed products of the form $M = R \rtimes \Gamma$ with R the hyperfinite factor.

It is interesting to note that Theorem 5.1 can be used to give a completely new proof of the by now classical result of Connes and Jones showing that property (T) factors cannot be embedded into the free group factor ([CJ]). Thus, rather than using Haagerup's property (i.e. "deformation by compact c.p. maps"), as the original proof does, this new proof uses a "deformation by automorphisms" of the free group factors.

6.7. Corollary. ([CJ]) *For every n , $2 \leq n \leq \infty$ the free group von Neumann algebra $L(\mathbb{F}_n)$*

contains no relatively rigid diffuse subalgebra.

Proof. If we write $L(\mathbb{F}_n)$ as $L(\mathbb{Z}) * L(\mathbb{Z}) * \dots * L(\mathbb{Z})$ and then apply recursively the first part of Theorem 5.1 and Corollary 3.2, it follows that a corner of $L(\mathbb{Z})$ contains a rigid diffuse von Neumann subalgebra, a contradiction. $\square$

7. AMALGAMATION OVER CARTAN SUBALGEBRAS: vNE/OE RIGIDITY RESULTS

In this section we apply Theorem 5.1 to study group measure space factors of the form $A \rtimes_{\sigma} \Gamma$ where Γ is a free product of groups $\Gamma = \Gamma_0 * \Gamma_1 * \dots$ and σ is a free ergodic m.p. action of Γ on $A = L^{\infty}(X, \mu)$, for (X, μ) a probability space. Such a factor can alternatively be viewed as a free product with amalgamation $M = M_0 *_A M_1 *_A \dots$, where $M_i = A \rtimes_{\sigma_i} \Gamma_i$, $\sigma_i = \sigma|_{\Gamma_i}$, with $A \subset M$, with the algebra of coefficients A of the crossed product $A \rtimes \Gamma$ now becoming the “core” of the amalgamated free product. It is this form that will allow us to use 5.1.

Following ([P6]), we regard an isomorphism of such group measure space factors as a *von Neumann equivalence* (vNE) of the corresponding actions $(\sigma, *_i \Gamma_i)$. Thus, the main result we prove in this section is a rigidity result showing that vNE of actions of free products of groups Γ_i satisfying some weak rigidity conditions (of property (T) type) entails the orbit equivalence (OE) of the actions σ , with component by component OE of the actions (σ_i, Γ_i) . Due to its analogy to similar statements on (amalgamated) free products of groups in Bass-Serre theory, we refer to this as *vNE Bass-Serre rigidity*. We note that when applied to isomorphisms of group measure space factors that come from OE of the actions, they give OE *Bass-Serre rigidity* results.

Since we study the group measure space factors $M = A \rtimes (\Gamma_0 * \Gamma_1 * \dots)$ as AFP factors $(A \rtimes \Gamma_0) *_A (A \rtimes \Gamma_1) *_A \dots$, it is worth noticing that if $M = M_0 *_A M_1 *_A \dots$ is an AFP factor coming from Cartan subalgebra inclusions $A = L^{\infty}(X, \mu) \subset M_i$, $i \geq 0$, then the AFP “core” A follows regular in M , but in general it may not be maximal abelian (thus not Cartan) in M . For instance, if A is a Cartan subalgebra of a II_1 factor N then A is not maximal abelian in $M = N *_A N$, because for any $u \in \mathcal{N}_N(A)$ with $E_A(u) = 0$ the element $u * u^{-1}$ is still perpendicular to A yet acts trivially on it. For more on general properties of AFP factors arising from Cartan inclusions we send the reader to ([Ko], [U1,2]).

In case $M_i = A \rtimes_{\sigma_i} \Gamma_i$, with each σ_i a free m.p. action, then there exists a unique m.p. action σ of $\Gamma = \Gamma_0 * \Gamma_1 * \dots$ on A such that $\sigma|_{\Gamma_i} = \sigma_i$, $\forall i$, and we still have the natural identification $M = A \rtimes_{\sigma} \Gamma = M_0 *_A M_1 *_A \dots$, as in the case σ a free action mentioned above. Then A is Cartan in M iff σ is a free action, i.e. iff σ_i are “freely independent” actions (in the obvious sense). For general equivalence relations (or Cartan subalgebras), the definition of “free independence” was formulated by Gaboriau ([G1]) and is reminded below. Recall that if B is a finite von Neumann algebra, p, q are non-zero projections in B and $\theta : pBp \rightarrow qBq$ is a $*$ -morphism, then θ is called *properly outer* if $b \in B$, $\theta(x)b = bx$, $\forall x \in B$, implies $b = 0$.

7.1. Definition ([G1]). Let $\{\mathcal{R}_i\}_{i \in I}$ be a family of countable, measurable, measure preserving equivalence relations on the same standard non atomic probability space (X, μ) (see

e.g. [FM]). We alternatively view each $\mathcal{R}_i$ as a pseudogroup of *local m.p. isomorphisms* $\phi : Y_1 \simeq Y_2$ with $Y_1, Y_2 \subset X$ measurable and the graph of ϕ contained in $\mathcal{R}_i$ ([Dy], [FM]). We say that $\{\mathcal{R}_i\}_i$ are *freely independent* if for any n and any properly outer local isomorphisms $\phi_j \in \mathcal{R}_{i_j}$, $1 \leq j \leq n$, $i_j \in I$, with $i_j \neq i_{j+1}$, $1 \leq j \leq n-1$, the product $\phi_1 \phi_2 \dots \phi_n$ is properly outer.

In the case when each of the equivalence relations $\mathcal{R}_i$ is generated by properly outer automorphisms, Definition 7.1 can be viewed as a particular case of the following:

7.1'. Definition. Let (B, τ) be a finite von Neumann algebra and $S_i \subset \text{Aut}(B, \tau)$, $i \in I$, a family of sets of τ -preserving automorphisms, with each $\theta \in S_i$ either properly outer or equal to id_B . We say that $\{S_i\}_i$ are *freely independent* if for any n and any $\theta_j \in S_{i_j} \setminus \{\text{id}_B\}$, $1 \leq j \leq n$, $i_j \in I$, with $i_1 \neq i_2 \neq \dots \neq i_n$, the product $\theta_1 \theta_2 \dots \theta_n$ is properly outer.

The next lemma translates the freeness conditions in definition 7.1 into the framework of operator algebras (see [U2]):

7.2. Lemma. Let (M_n, τ_n) , $n \geq 1$, be finite von Neumann algebras with a common Cartan subalgebra $A \subset M_n$, such that $\tau_n|_A = \tau_m|_A$, $\forall n, m$. Then $A \subset M_1 *_A M_2 *_A \dots$ is a Cartan subalgebra if and only if the equivalence relations $\mathcal{R}_n = \mathcal{R}_{A \subset M_n}$, $n \geq 1$, are freely independent.

Proof. Since A is clearly regular in $M = M_1 *_A M_2 *_A \dots$, all we need to prove is that $A \subset M$ is maximal abelian iff $\{\mathcal{R}_n\}_n$ are freely independent. But this is trivial by the definitions of freeness and respectively of the amalgamated free product over A . $\square$

The next result, essentially due to Tornquist ([To]), shows that a sequence of actions of countable groups (or merely countable m.p. equivalence relations) can be made “freely independent” by conjugating each one of them with a suitable m.p. automorphism. We include a proof, based on Lemma A.1 in the Appendix, for the reader’s convenience.

7.3. Proposition. 1°. Let (X, μ) be a standard non atomic probability space and $\sigma_n : G_n \rightarrow \text{Aut}(X, \mu)$ be free m.p. actions of discrete countable groups G_n , $n \geq 1$. Then there exists a free m.p. action σ of $G = *_n G_n$ on (X, μ) such that $\sigma|_{G_n}$ is conjugate to σ_n , $\forall n \geq 1$. More generally, if $\{\mathcal{R}_n\}_n$ are standard equivalence relations on (X, μ) then there exists an equivalence relation $\mathcal{R}$ on (X, μ) generated by a family of freely independent sub equivalence relations $\mathcal{R}'_n \subset \mathcal{R}$, $n \geq 1$, such that $\mathcal{R}_n \simeq \mathcal{R}'_n$, $\forall n$.

2°. Let (M_n, τ_n) be countably generated finite von Neumann algebras with a common diffuse Cartan subalgebra $A \subset M_n$, $n \geq 0$, such that $\tau_n|_A = \tau_m|_A$, $\forall n, m$. Then there exist Cartan subalgebra inclusions $\{A \subset N_n\}_n$ such that $(A \subset N_n) \simeq (A \subset M_n)$, $\forall n$, and such that A is a Cartan subalgebra in $N_0 *_A N_1 *_A N_2 \dots$.

Proof. 1°. This is an immediate application by induction of Lemma A.1, once we notice that any $\mathcal{R}_n$ can be extended to a countable m.p. equivalence relation $\mathcal{S}_n$ on (X, μ) which is generated by countably many properly outer m.p. automorphisms.

2°. By Lemma 7.2, we only need to make the equivalence relations $\mathcal{R}_{A \subset M_n}$ freely independent, so the first part applies. $\square$

We also notice the following general “compression formula” for restrictions of “free products of equivalence relations” $\mathcal{R} = *_{i=1}^n \mathcal{R}_i$, i.e. for relations $\mathcal{R}$ that are generated by freely independent sub-equivalence relations $\mathcal{R}_i \subset \mathcal{R}, 1 \leq i \leq n$.

7.4. Proposition. 1°. Let $M_i, 1 \leq i \leq n$, be II_1 factors, for some $2 \leq n \leq \infty$, with a common Cartan subalgebra A and assume $A \subset M = M_1 *_A M_2 \dots *_A M_n$ is Cartan. If $p \in A$ is a projection of trace $1/m$ for some integer $m \geq 1$ then the Cartan subalgebra inclusion $Ap \subset pMp$ is naturally isomorphic to $Ap \subset M_0 *_A pM_1p *_A p \dots *_A pM_np$, where $(Ap \subset M_0) = (Ap \rtimes \mathbb{F}_{(n-1)(m-1)})$ for some free action of the free group $\mathbb{F}_{(n-1)(m-1)}$ on Ap .

2°. Let $\mathcal{R}_1, \dots, \mathcal{R}_n$ be freely independent, countable, ergodic, m.p. equivalence relations on the same standard probability space (X, μ) and denote by $\mathcal{R}$ the equivalence relation they generate. If $Y \subset X$ is a subset of measure $1/m$ then the restriction $\mathcal{R}^Y$ of $\mathcal{R}$ to Y is generated by the freely independent ergodic sub-equivalence relations $\mathcal{R}_i^Y, 1 \leq i \leq n$, and $\mathcal{R}_0$, where $\mathcal{R}_i^Y$ is the restriction of $\mathcal{R}_i$ to Y and $\mathcal{R}_0$ is generated by a free m.p. action of a free group with $(n-1)(m-1)$ generators $\mathbb{F}_{(n-1)(m-1)}$ on Y .

Proof. It is clearly sufficient to prove part 1°. By the representation of AFP algebras (1.1.1), the von Neumann algebra generated by $pM_ip, 1 \leq i \leq n$, in pMp is isomorphic to the AFP algebra $pM_1p *_A p \dots *_A pM_np$. On the other hand, since $\tau(p) = 1/m$ and each M_i is a factor, by Dye’s Theorem ([D]) there exist matrix units $\{e_i^{jk}\}_{1 \leq j,k \leq m}$ in the normalizing groupoid $\mathcal{GN}_{M_i}(A)$ of $A \in M_i$ such that $e_i^{11} = p$ and $e_i^{jj} = e_{i'}^{jj}, \forall 1 \leq i, i' \leq n$ and $\forall 1 \leq j \leq m$. Denote by $u_i^j = e_1^{1j} e_i^{j1} \in pMp, 2 \leq i \leq n, 2 \leq j \leq m$, and notice that there are $(n-1)(m-1)$ many such unitary elements. The expansion (1.1.1) of $M_1 *_A \dots *_A M_n$ implies that $\{u_i^j\}_{i,j}$ are the generators of a free group $\mathbb{F}_{(n-1)(m-1)}$, all of whose elements $\neq 1$ are perpendicular to Ap . Since $A \subset M$ is Cartan, this implies the action implemented by $\mathbb{F}_{(n-1)(m-1)}$ on Ap is free. Moreover, if we denote $M_0 = Ap \vee \{u_i^j\}_{i,j}'' \simeq Ap \rtimes \mathbb{F}_{(n-1)(m-1)}$ then it is immediate to check that if $v = v_{i_1} v_{i_2} \dots v_{i_k} \in pMp$ is an “alteranting word”, with v_{i_l} in the normalizing groupoid of Ap in $pM_{i_l}p, \forall l$, and $i_1 \neq i_2 \neq \dots \neq i_k$ in $\{0, 1, \dots, n\}$, $E_{Ap}(v_{i_l}) = 0, \forall l$, then v has expectation 0 on Ap as well, $E_{Ap}(v) = 0$. Thus, if we denote by $N \subset pMp$ the von Neumann algebra generated by $pM_ip, 0 \leq i \leq n$, then $(Ap \subset N) = (Ap \subset M_0 *_A pM_1p *_A p \dots *_A pM_np)$.

Finally, since $e_1^{j1} u_i^j = e_i^{j1}$, we have that M is generated by $N = pNp$ and the matrix unit $\{e_1^{jk}\}_{j,k}$. This also implies that pMp is generated by $\vee_{i=0}^n pM_ip$. Altogether, this shows that $(Ap \subset pMp) = (Ap \subset N) = (Ap \subset M_0 *_A pM_1p *_A p \dots *_A pM_np)$. $\square$

The above result shows in particular that if $\mathbb{F}_n \curvearrowright X$ is a free m.p. action on the probability space with restrictions to each of the generators of $\mathbb{F}_n$ acting ergodically on X then the amplification of the corresponding orbit equivalence relation $\mathcal{R}_{\mathbb{F}_n}$ by $1/m$ is an equivalence relation that can be implemented by a free ergodic m.p. action of a free group with $m(n-1) + 1$ generators, thus recovering the “compression formula” $(\mathcal{R}_{\mathbb{F}_n})^{1/m} = \mathcal{R}_{\mathbb{F}_{m(n-1)+1}}$ in ([Hj]). On the other hand, Proposition 7.4 can be viewed as an AFP version of the “compression formula” for plain free product factors in ([DyR]).

The statement 7.5 below will require the following:

7.5.0. Notation. Let $\{\Gamma_{i,j}\}_{j=0}^{n_i}$ be discrete countable groups, $1 \leq n_i \leq \infty$, $i = 1, 2$. Denote $G_i = \Gamma_{i,0} * \Gamma_{i,1} * \dots * \Gamma_{i,n_i}$, $i = 1, 2$. Let $\sigma_i : G_i \rightarrow \text{Aut}(X_i, \mu_i)$ be a free, m.p. action on a standard probability space (X_i, μ_i) . Denote $A_i = L^\infty(X_i, \mu_i)$, $M_i = A_i \rtimes_{\sigma_i} G_i$ and $M_{ij} = A_i \rtimes_{\sigma_{ij}} \Gamma_{ij}$, where $\sigma_{ij} = (\sigma_i)|_{\Gamma_{ij}}$, $\forall n_i \geq j \geq 0, i = 1, 2$.

The general result we prove shows that, under suitable weak rigidity conditions on the groups Γ_{ij} , an isomorphism between the factors M_1, M_2 must take each of the “component inclusions” ($A_i \subset M_{ij}$) onto each other, modulo some permutation of indices and unitary conjugacy. Since the weak rigidity assumption on the Γ_{ij} ’s is somewhat technical, we display the conditions separately and give right away a list of examples when they are satisfied:

7.5.1. Assumption. Γ_{10}, Γ_{20} have Haagerup property, $i = 1, 2$, and if both Γ_{10}, Γ_{20} are finite then at least one of the n_i must be ≥ 2 . For each $j \geq 1$, $i = 1, 2$, Γ_{ij} contains a subgroup H_{ij} with the following properties:

- (a). H_{ij} is non virtually abelian and the pair (Γ_{ij}, H_{ij}) has the relative property (T) ([M]).
- (b). The normalizer N_{ij} of H_{ij} in Γ_{ij} is ICC in Γ_{ij} (i.e. $|\{hgh^{-1} \mid h \in N_{ij}\}| = \infty$, $\forall g \in \Gamma_{ij} \setminus \{e\}$) and σ_{ij} is ergodic on N_{ij} .
- (c). For any proper intermediate subgroup $N_{ij} \subset N'_{ij} \subset \Gamma_{ij}$ there exists $g \in \Gamma_{ij} \setminus N'_{ij}$ such that $g(N'_{ij})g^{-1} \cap N'_{ij}$ is non virtually abelian.

Note that condition (c) above on the inclusion $N_{ij} \subset \Gamma_{ij}$ is similar to the *wq-normal* condition in ([P6,8]). It is equivalent to the existence of a well ordered, strictly increasing family of intermediate subgroups $\{G_l \mid 0 \leq l \leq L\}$ such that $G_0 = N_{ij}$, $G_L = \Gamma_{ij}$ and $G_{k+1} = \{g \in \Gamma_{ij} \mid gG_kg^{-1} \cap G_k \text{ non virtually abelian}\}$, $\forall k$.

If $\Gamma_{ij}, j \geq 1$, is ICC and has a normal, non virtually abelian subgroup with the relative property (T) then conditions (a), (b), (c) are trivially satisfied.

Related to conditions (a) and (b), note that if a non virtually abelian group H is normal in an ICC group G then $L(H)$ has no type I summand, i.e. it is of type II_1 . Indeed, because G acts ergodically on the center of $L(H)$, so if $L(H)$ has non-zero type I part then it is homogeneous of type I_n for some $2 \leq n < \infty$, contradicting ([Ka]).

7.5. Theorem (vNE Bass-Serre rigidity). *With the notations 7.5.0 and assumptions 7.5.1, if $\theta : M_1 \simeq M_2^t$, for some $t > 0$, then $n_1 = n_2$ and there exists a permutation π of indices $j \geq 1$ and unitaries $u_j \in M_2^t$ such that $\text{Ad}(u_j)(\theta(M_{1,j})) = M_{2,\pi(j)}^t$, $\text{Ad}(u_j)(\theta(A_1)) = A_2^t$, $\forall j \geq 1$. In particular, $\mathcal{R}_{\sigma_1} \simeq \mathcal{R}_{\sigma_2}^t$ and $\mathcal{R}_{\sigma_{1,j}} \simeq \mathcal{R}_{\sigma_{2,\pi(j)}}^t$, $\forall j \geq 1$. Also, if $\Gamma_{10} = \Gamma_{20} = 1$, $2 \leq n_1 < \infty$ then the existence of such an isomorphism forces $t = 1$.*

Proof. We denote by $Q_{ij} \subset M_{ij}$ the “rigid part” of M_{ij} , i.e. $Q_{ij} = L(H_{ij})$ where $H_{ij} \subset \Gamma_{ij}$ is a subgroup satisfying properties (a), (b), (c) in 7.5.1. Also, we denote by P_{ij} the von Neumann algebra generated by the normalizer of Q_{ij} in M_{ij} . Thus, $P_{ij} \supset L(N_{ij})$, where N_{ij} is the normalizer of H_{ij} in Γ_{ij} in (b). Notice that conditions 7.5.1 imply that $P'_{ij} \cap M_{ij} = \mathbb{C}$ so that by 1.2.1 we also have $P'_{ij} \cap M_i = \mathbb{C}$, $\forall j \geq 1, i = 1, 2$. Also, note that Q_{ij} is relatively rigid in M_{ij} , thus in M_i .

Assume first that $t \leq 1$. For simplicity, denote $A_2 = A$, $M_2 = M$, $G_2 = G$, $\Gamma_{2j} = \Gamma_j$ and let $\{u_g\}_{g \in G} \subset M$ denote the canonical unitaries. Let $q \in A$ be so that $\tau(q) = t$ and $\theta(1) = q$. Fix $i \geq 1$. Since $Q_{1i} \subset M_1$ is a rigid inclusion, $Q = \theta(Q_{1i}) \subset qMq$ is a rigid inclusion ([P5]).

Let us show that no corner of Q can be embedded into M_{20} inside M . Assume by contradiction that there exist non-zero $q_0 \in \mathcal{P}(Q)$, $p_0 \in \mathcal{P}(M_{20})$, a unital isomorphism ψ of q_0Qq_0 into $p_0M_{20}p_0$ and a non-zero partial isometry $v \in M_{20}$ such that $v^*v \in (q_0Qq_0)' \cap q_0Mq_0$, $vv^* \in \psi(q_0Qq_0)' \cap p_0Mp_0$ and $vy = \psi(y)v$, $\forall y \in q_0Qq_0$. Since Q is type II, $\psi(q_0Qq_0) \subset M_{20}$ is type II, so no corner of $\psi(q_0Qq_0)$ can be embedded into A . By 1.2.1, this implies $\psi(q_0Qq_0)' \cap M \subset M_{20}$. Thus $vQv^* = v(q_0Qq_0)v^* \subset M_{20}$. But Γ_{20} has Haagerup property, so by ([P5]) there exist unital trace preserving A -bimodular c.p. maps ϕ_n on M_{20} such that $\phi_n \rightarrow id_{M_{20}}$ and ϕ_n compact relative to A . Then $\phi_n * id \rightarrow id_M$. But by ([P5]), $vQv^* \subset vv^*Mvv^*$ is a rigid inclusion. By ([P5]) again this implies $\lim_n \|(\phi_n * id)(x) - x\|_2 = 0$ uniformly for $x \in (vQv^*)_1$. Since $(\phi_n * id)(x) = \phi_n(x)$ for $x \in M_{20} \supset vQv^*$, this implies the maps ϕ_n , which are A -bimodular and compact relative to A , tend uniformly to the identity on the unit ball of the type II₁ algebra vQv^* . By ([P5]), this implies a corner of vQv^* can be embedded into A inside M_{20} , a contradiction.

Since no corner of Q can be embedded into M_{20} inside M we can apply Theorem 5.1. Thus, there exist $j = j(i) \geq 1$, a non-zero projection $q' \in Q' \cap qMq$ and a unitary element $u \in M$ such that $uQq'u^* \subset qM_{2,j}q = q(A \rtimes \Gamma_{2,j})q$. Since $P = \theta(P_{1i})$ is generated by the normalizer of Q , we have $q' \in P$ and by (3.5.1° in [P6]), $Qq' \subset q'Pq'$ is quasi-regular. By Theorem 1.2.1 it follows that $uq'Pq'u^* \subset qM_{2,j}q$. Since P is a factor and a corner of it is contained in $qM_{2,j}q$, it follows that u can be suitably modified so that to satisfy $uPu^* \subset qM_{2,j}q$. If $v \in qMq$ is such that $P_v = vPv^* \cap P$ is type II₁, then $P_v \subset qM_{2,j}q$ and no corner of P_v can be embedded into the abelian algebra A . Since $(P_v)v = vP \subset vM_{2,j}$, by Theorem 1.2.1 it follows that $v \in M_{2,j}$. By applying this recursively, by 7.5.1 (c) it follows that $\theta(L(\Gamma_{1i})) \subset qM_{2,j}q$.

Thus, if $\{u_{1,h} \mid h \in G_1\}$ denotes the canonical unitaries in $M_1 = A_1 \rtimes G_1$, then $v_h = \theta(u_{1,h}) \in qMq$, $h \in G_1$, are in the normalizer of $\theta(A_1)$ in qMq . Thus, the unitary elements $\{uv_hu^* \mid h \in \Gamma_{1i}\} \subset qM_{2,j}q$ normalize $u\theta(A_1)u^*$ and they generate a II₁ von Neumann algebra $N \subset qM_{2,j}q$. By Proposition 1.4.1, this implies $u\theta(A_1)u^* \subset M_{2,j}$ and $w(u\theta(A_1)u^*)w^* = Aq$ for some $w \in \mathcal{U}(qM_{i,j}q)$. Taking $v = wu$, it follows that $v(\theta(L(\Gamma_{1,i})))v^* \subset M_{2,j}$ and $v\theta(A_1)v^* \subset M_{2,j}$, thus $v\theta(M_{1,i})v^* \subset M_{2,j}$.

This shows that if $t \leq 1$ then $\forall i \geq 1$ there exists $j = j(i) \geq 1$ (unique by Theorem 1.2.1) and a unitary element $v \in M_2^t$ such that $\text{Ad}(v)\theta(M_{1,i}) = M_{2,j}^t$ and $\text{Ad}(v)(\theta(A_1)) = A_2^t$.

Let us now consider the case $t > 1$. Let $n \geq t$ be an integer and note that if we let $B = M_{n \times n}(A_2) \subset M_{n \times n}(M_2) = M$, $G = G_2$ and extend σ_2 to the action σ which acts trivially on $M_{n \times n}(\mathbb{C}) \subset M_{n \times n}(M_2) = M$, then $M = B \rtimes_\sigma G_2$. Let $q \in A = D_n \otimes A_2$ be a projection of trace t/n .

The hypothesis then states that $\theta : M_1 \simeq qMq$ is an onto isomorphism. Fix $i \geq 1$ and let $Q = \theta(Q_{1i}) \subset qMq$, $P = \theta(P_{1i}) \subset qMq$. As in the case $t \leq 1$, it follows that there exist $j = j(i) \geq 1$ and a unitary element u in $qMq = M_2^t$ such that $uPu^* \subset q(B \rtimes \Gamma_{2,j})q = M_{2,j}^t$.

In particular, $\{\theta(u_{1,h}) \mid h \in \Gamma_{1,i}\} \subset qM_{2,j}q$ and they normalize $u\theta(A_1)u^*$. By applying Theorem 1.4.1 to $A_0 = u\theta(A_1)u^* \subset qBq$, it follows that $u(\theta(A_1))u^* \subset q(B \rtimes \Gamma_{2,j})q$ as well, so $\text{Ad}(u)(\theta(M_{1,i})) \subset M_{2,j}^t$. Since $\text{Ad}(u)(\theta(A_1))$ is regular in $M_2^t = qMq$, by Corollary 1.2.4 and (A.1 in [P5]) it follows that there exists a unitary in $M_{2,j}$ that conjugates $\theta(A_1)$ onto A_2^t .

Since we have dealt with both cases $t \geq 1$ and $t \leq 1$, we can apply the above equally well to θ and θ^{-1} , to obtain the following: $\forall n_1 \geq i \geq 1, n_2 \geq j \geq 1$, there exist unitaries $u_i \in M_2^t, v_j \in M_1$ and indices $j(i) \in \{1, 2, \dots, n_2\}, i(j) \in \{1, 2, \dots, n_1\}$ such that

$$\theta(A_1 \subset M_{1,i}) \subset u_i(A_2 \subset M_{2,j(i)})^t u_i^*, 1 \leq i \leq n_1,$$

$$\theta^{-1}((A_2 \subset M_{2,j})^t) \subset v_j(A_1 \subset M_{1,i(j)})v_j^*, 1 \leq j \leq n_2,$$

so altogether

$$\theta(M_{1,k}) \subset \text{Ad}(u_k \theta(v_{j(k)}))(M_{1,i(j(k))}), 1 \leq k \leq n_1.$$

which by Theorem 1.1 entails $i(j(k)) = k, \forall k$. Similarly $j(i(k)) = k, \forall k$. Thus, $n_1 = n_2 = n$ and j defines a permutation π of the set of indices $1 \leq i \leq n$.

To prove the last part note that the equivalence $\mathcal{R}_{\sigma_1} \simeq \mathcal{R}_{\sigma_2}^t$ and ([G1]) imply

$$\sum_{j=1}^n \beta_1(\Gamma_{1j}) + (n-1) = \beta_1(\mathcal{R}_{\sigma_1}) = \beta_1(\mathcal{R}_{\sigma_2})/t = (\sum_{j=1}^n \beta_1(\Gamma_{2j}) + (n-1))/t.$$

On the other hand, by the equivalence $\mathcal{R}_{\sigma_{1j}} \simeq \mathcal{R}_{\sigma_{2,\pi(j)}}^t$ we get $\beta_1(\Gamma_{1j}) = \beta_1(\Gamma_{2,\pi(j)})/t, \forall j \geq 1$ while by ([BeVa]) and assumptions 7.5.1 all Γ_{1j} 's have 0 first ℓ^2 -Betti number, $\beta_1(\Gamma_{1j})$. (Indeed, this follows easily from Corollary 4 in [BeVa] and the argument in the proof of 2.4 in [P8].) Altogether we get $n-1 = (n-1)/t$, implying that $t = 1$. $\square$.

Before stating specific OE applications, recall from ([Fu1]) that two groups Γ, Λ are said to be *measure equivalent* (ME) with *dilation constant* $t > 0$ if there exists free, m.p. actions $(\sigma, \Gamma), (\theta, \Lambda)$ such that $\mathcal{R}_\sigma \simeq \mathcal{R}_\theta^t$. We'll use the notation $\Gamma \sim_{OE_t} \Lambda$ to denote this property. It was recently proved in ([G2]) that if $\Gamma_i \sim_{OE_1} \Lambda_i, \forall i \geq 0$, then $*_j \Gamma_j \sim_{OE_1} *_j \Lambda_j$. Note that an alternative proof of this fact follows from 7.3.1° (see also the comment 2.27 in [MS]).

The OE rigidity result below, of Bass-Serre type, can alternatively be viewed as a converse to Gaboriau's above ME result, for free products of w-rigid ICC groups. Note however that we need the actions involved to be “separately ergodic” (not assumed in [G2]).

7.6. Corollary (OE Bass-Serre rigidity). *Let Γ_0, Λ_0 be Haagerup groups and $\Gamma_i, \Lambda_j, 1 \leq i \leq n \leq \infty, 1 \leq j \leq m \leq \infty$, be ICC groups having normal, non virtually abelian subgroups with the relative property (T). Let σ (resp. θ) be a free ergodic m.p. action of $\Gamma = \Gamma_0 * \Gamma_1 * \dots$ (resp. $\Lambda = \Lambda_0 * \Lambda_1 * \dots$) on the probability space such that $\sigma_j = \sigma|_{\Gamma_j}$ (resp. $\theta_j = \theta|_{\Lambda_j}$) is egodic $\forall j \geq 1$. If $\mathcal{R}_{\sigma,\Gamma} \simeq \mathcal{R}_{\theta,\Lambda}^t$ then $n = m$ and there exists a permutation π of the set of indices ≥ 1 such that $\mathcal{R}_{\sigma_i,\Gamma_i} \simeq \mathcal{R}_{\theta_{\pi(i)},\Lambda_{\pi(i)}}^t, \forall i \geq 1$.*

The condition on the groups $\Gamma_i, \Lambda_j, i, j \geq 1$, in 7.6 can be weakened by using the full generality of Theorem 7.5:

7.6'. Corollary. *The same statement as in 7.6 holds true if we assume each $(\sigma_{|\Gamma_i}, \Gamma_i)$ (resp. $(\theta_{|\Lambda_i}, \Lambda_i)$), $i \geq 1$, satisfies conditions 7.5.1(a), (b), (c).*

7.7. Corollary. *Let $\Gamma_i, 0 \leq i \leq n$, $\Gamma = \Gamma_0 * \Gamma_1 * \dots$, σ be as in 7.6 or 7.6'. Assume $\text{Out}(\mathcal{R}_{\sigma_1, \Gamma_1}) = \{1\}$ and (σ_1, Γ_1) is not orbit equivalent to (σ_i, Γ_i) for any $i \neq 1$. Then $\text{Out}(\mathcal{R}_{\sigma, \Gamma}) = \{1\}$ and $\text{Out}(A \rtimes_{\sigma} \Gamma) = H^1(\sigma, \Gamma)$. Also, if n is finite and either all Γ_i are finitely generated or there exists $i \geq 1$ with $\beta_n(\Gamma_i) \neq 0, \infty$, then $\mathcal{F}(A \rtimes_{\sigma} \Gamma) = \{1\}$.*

Proof. The first part is trivial by Theorem 7.5 while the last part follows from ([G1]). $\square$

Outer automorphism groups of equivalence relations are usually hard to calculate and there are only a few special families of group-actions (σ_1, Γ_1) for which one knows that $\text{Out}(\mathcal{R}_{\sigma_1}) = \{1\}$ (cf. [Ge2], [Fu2], [MS]). Similarly for the 1-cohomology group H^1 , where the only known calculations are in ([Ge1], [PSa], [P8]). We remind below some examples from ([Ge2], [Fu2], [MS]) where both calculations can be made. We add a new construction of examples, in 7.8.3 below, which uses the Monod-Shalom OE rigidity theorem to calculate Out and ([P8]) to calculate H^1 .

7.8.1. Example (Gefer [Ge2], Furman [Fu2]). Take Γ_1 a lattice in $SO(p, q)$, with $p > q \geq 2$ and notice that Γ_1 has $\text{rk}_{\mathbb{R}} \geq 2$ (thus has property T) and admits a dense embedding into the compact Lie group $SO(n)$, where $n = p + q$. Let σ_1 be the action by left translation of Γ_1 on the homogeneous space $SO(n)/SO(n-1)$. Then $\text{Out}(\mathcal{R}_{\sigma_1}) = \{1\}$.

7.8.2. Example (Monod-Shalom [MS]). Let $\mathcal{G} = SO(n), n \geq 5$, and Λ_0 an ICC, torsion free, Kazhdan group which admits a dense embedding into $\mathcal{G}$ and has no outer automorphisms (such groups exists for any $n \geq 5$, for instance lattices in $SO(p, q)$ with $p > q \geq 2$ and $n = p + q$ as in 7.8.1). Let K be any torsion free group embeddable into $\mathcal{G}$ and $K_0 \subset K$ a non-trivial subgroup such that K_0 is not isomorphic to K (for instance, $K_0 = \mathbb{F}_r \subset \mathbb{F}_s = K$, for some $s > r \geq 1$). Let $\Gamma_1 = (\Lambda_0 * K) \times (\Lambda_0 * K_0)$ and note that any automorphism of the group Γ_1 is inner on $\Lambda_0 \times \Lambda_0$ (by Kurosh or Bass-Serre). Let σ_1 be the action of $\Gamma_1 = (\Lambda_0 * K) \times (\Lambda_0 * K_0)$ on $\mathcal{G}$ by left-right translation. Notice that this action is free ergodic on $\Lambda_0 \times \Lambda_0$ and that by ([MS]) one can choose the embedding $K \subset \mathcal{G}$ such that σ_1 is free on Γ_1 (by considering all embeddings gKg^{-1} , $g \in \mathcal{G}$, and using a Baire category argument). Then Γ_1 is in the class $\mathcal{C}_{geom}$ of Monod-Shalom and $\text{Out}(\mathcal{R}_{\sigma_1}) = \{1\}$.

7.8.3. Example. Let this time Λ_0 be any torsion free ICC group with only inner automorphisms and which cannot be decomposed as $\mathbb{Z} * \Lambda'_0$ (note that if Λ_0 is w-rigid then it does have this latter “free in-decomposability” property). Let K be any torsion free group with $K_0 \subset K$ a non-trivial subgroup such that K_0 is not isomorphic to K . Let $\Lambda = \Lambda_0 * K$ and $\Gamma_1 = (\Lambda_0 * K) \times (\Lambda_0 * K_0) \subset \Lambda \times \Lambda$. Denote by σ_1 the action of Γ_1 on the product probability space $(X, \mu) = \prod_{g \in \Lambda} (X_0, \mu_0)_g$ by left-right (double) Bernoulli shifts. Then $\text{Out}(\mathcal{R}_{\sigma_1}) = \{1\}$. Indeed, noticing that $\Lambda_0 * K, \Lambda_0 * K_0 \in \mathcal{C}_{geom}$ and that σ_1 is separately ergodic (on $\Lambda_0 * K$ and $\Lambda_0 * K_0$) it follows from ([MS]) that any automorphism $\theta \in \text{Aut}(\mathcal{R}_{\sigma_1})$ is an inner perturbation of a conjugacy $\sigma_1 \sim \sigma_1 \circ \gamma$ with respect to some $\gamma \in \text{Aut}(\Gamma_1)$. But Γ_1 has only inner automorphisms (again by Kurosh or Bass-Serre). Thus, any such θ is an inner perturbation of an automorphism of the probability space that commutes with $\sigma_1(\Gamma_1)$.

But this commutant is trivial if for instance (X_0, μ_0) is atomic with un-equal weights, as shown by the following:

7.9. Proposition. *Let Λ be a countable discrete group with two subgroups $\Lambda_1, \Lambda_2 \subset \Lambda$ such that $\Lambda_0 = \Lambda_1 \cap \Lambda_2$ satisfies $|\{hgh^{-1} \mid h \in \Lambda_0\}| = \infty, \forall g \in \Lambda, g \neq e$. Let (B_0, τ_0) be a finite von Neumann algebra and denote $(B, \tau) = \Pi_{g \in \Lambda}(B_0, \tau_0)_g$. Let σ be the action of $\Lambda_1 \times \Lambda_2$ on (B, τ) given by $\sigma_{h_1, h_2}((x_g)_g) = (x'_g)_g$, where $x'_g = x_{h_1^{-1}gh_2}$, $\forall g \in \Lambda, h_i \in \Lambda_i, i = 1, 2$. Then σ is a free, separately mixing action of $\Lambda_1 \times \Lambda_2$ on (B, τ) and we have:*

- 1°. *If $\theta \in \text{Aut}(B, \tau)$ commutes with $\sigma(\Lambda_1 \times \Lambda_2)$ then there exists a unique $\theta_0 \in \text{Aut}(B_0, \tau_0)$ such that θ is the product action given by θ_0 , i.e. $\theta = \Pi_g(\theta_0)_g$.*
- 2°. *Any $\delta \in \text{Aut}(\Lambda)$ satisfying $\delta(\Lambda_i) = \Lambda_i, i = 1, 2$, implements an automorphism $\Delta = \Delta(\delta) \in \text{Aut}(B, \tau)$, by $\Delta((b_g)_g) = (\delta(b_g))_g$, which satisfies $\Delta\sigma\Delta^{-1} = \sigma \circ \delta$.*
- 3°. *If $(B_0, \tau_0) = (L^\infty(X_0, \mu_0), \int \cdot d\nu_0)$ for some atomic probability space (X_0, μ_0) , then the commutant of $\sigma(\Lambda_1 \times \Lambda_2)$ in $\text{Aut}(X, \mu)$ is equal to $\text{Aut}(X_0, \mu_0) = \text{Aut}(B_0, \tau_0)$. Moreover, if σ is conjugate to another double Bernoulli shift σ' with atomic base space (X'_0, μ'_0) , then $(X_0, \mu_0) \simeq (X'_0, \mu'_0)$.*

Proof. Part 2° is trivial. To prove part 1°, it is sufficient to show that any θ commuting with σ must take the subalgebra $B_0^e = \dots 1 \otimes (B_0)_e \otimes 1 \dots$ of B into itself. Indeed, because if we denote $\theta_0 = \theta|_{B_0^e}$ and regard it as an automorphism of B_0 then $\theta \circ (\Pi_g(\theta_0)_g)^{-1}$ still commutes with $\sigma(\Lambda_1 \times \Lambda_2)$ and it acts as the identity on B_0^e , thus on $\sigma(g_1, g_2)(B_0^e), \forall g_1, g_2 \in \Lambda_0$. Since $\Lambda_1 \Lambda_2 = \Lambda$, the latter generate all B , thus $\theta = \Pi_g(\theta_0)_g$.

To show that θ leaves B_0^e globally invariant, it is sufficient to show that the fixed point algebra $\{b \in B \mid \sigma(g, g)(b) = b, \forall g \in \Lambda_0\}$ coincides with B_0^e . This in turn follows trivially from the fact that for any finite subset $F \subset \Lambda \setminus \{e\}$ there exists $g \in \Lambda_0$ such that $gFg^{-1} \cap F = \emptyset$. To see that this latter property holds true, note that if some finite set $\emptyset \neq F \subset \Lambda \setminus \{e\}$ would satisfy $|gFg^{-1} \cap F| \geq 1, \forall g \in \Lambda_0$, then the “left-right” representation $\pi(g)(f) = \lambda(g)\rho(g)(f)$ on $\ell^2(\Lambda)$ would satisfy $\langle \pi(g)(\chi_F), \chi_F \rangle \geq |F|^{-1}$ for all $g \in \Lambda_0$. Taking the element f of minimal Hilbert norm in $\overline{\text{co}}^w\{\pi(g)(\chi_F) \mid g \in \Lambda_0\} \subset \ell^2(\Lambda)$, it follows that $f \geq 0, f \neq 0$ (because $\langle f, \chi_F \rangle \geq |F|^{-1}$) and $\pi(g)(f) = f, \forall g \in \Lambda$. But then any appropriate “level set” K for f will be finite, non-empty and will satisfy $\pi(g)(\chi_K) = \chi_K, \forall g \in \Lambda_0$, i.e. $gKg^{-1} = K, \forall g \in \Lambda_0$, implying that Λ has elements with finite conjugacy class, a contradiction.

The first part of 3° is trivial by 2°. Then to see that conjugacy of double Bernoulli shifts entails isomorphism of the base spaces, note that σ conjugate to σ' implies that all “diagonal” actions $\sigma \otimes \sigma'', \sigma' \otimes \sigma''$ must also be conjugate, $\forall \sigma''$, thus having isomorphic commutants in Aut . Taking σ'' to be itself a double Bernoulli Γ_1 -action of base (X''_0, μ''_0) , it follows that $\text{Aut}((X_0, \mu_0) \times (X''_0, \mu''_0)) \simeq \text{Aut}((X'_0, \mu'_0) \times (X''_0, \mu''_0)), \forall (X''_0, \mu''_0)$, which easily implies the result. $\square$

7.10.0. Notation. Denote by $w\mathcal{T}_2$ the class of groups G that have a non virtually abelian subgroup $H_0 \subset G$ such that: (G, H_0) is a property (T) pair; the normalizer H of H_0 in G satisfies $|\{hgh^{-1} \mid h \in H\}| = \infty, \forall g \in G \setminus \{e\}$; the wq-normalizer of H in G generates G .

Note that any group in $w\mathcal{T}_2$ is ICC and that if G is ICC and has a normal, non abelian, relatively rigid subgroup then $G \in w\mathcal{T}_2$. Thus, any group of the form $G = H_0 \times K$ with H_0 ICC Kazhdan and K either ICC or equal to 1 is w-rigid and thus in $w\mathcal{T}_2$. Also, if $G \in w\mathcal{T}_2$ then $(G * K_0) \times K \in w\mathcal{T}_2$ for any ICC group K and any arbitrary group K_0 .

7.10. Corollary. *Let Γ_0 be a Haagerup group and $\Gamma_i \in w\mathcal{T}_2$, for $1 \leq i \leq n$, where $1 \leq n \leq \infty$. Assume Γ_1 is as in 7.8.1, 7.8.2, or 7.8.3. Then $\Gamma = \Gamma_0 * \Gamma_1 * \dots$ has a free ergodic m.p. action σ with $\text{Out}(\mathcal{R}_\sigma) = \{1\}$. Moreover, we have:*

1°. *If Γ_1 is as in 7.8.2 or 7.8.3, then there exist uncountably many non stably orbit equivalent actions σ of Γ with $\text{Out}(\mathcal{R}_\sigma) = \{1\}$ and $\mathcal{F}(\mathcal{R}_\sigma) = \{1\}$.*

2°. *If in addition Γ_0 is a product of amenable groups then given any discrete countable abelian group K , the uncountable family of actions σ in 1° can be taken to satisfy $H^1(\sigma, \Gamma) = \mathbb{G}_0^{n-1} \times \mathbb{G} \times \prod_{j \geq 1} \text{Char}(\Gamma_j) \times K^{n-1}$, where $\mathbb{G}$ is the polish group $\mathcal{U}(L^\infty(\mathbb{T}, \lambda))$ and $\mathbb{G}_0 = \mathbb{G}/\mathbb{T}$.*

Proof. By 7.3, we can take the free m.p. action σ of Γ on $A = L^\infty(X, \mu)$ so that for each $i \neq 1$, $\sigma_i = \sigma|_{\Gamma_i}$ is a (left) Bernoulli Γ_i -action, or a quotient of it as in ([P8]), and of the form 7.8.1–7.8.3 for $i = 1$. Notice that in example 7.7.1 the group Γ_1 has property (T) and is ICC, thus $\Gamma_1 \in w\mathcal{T}_2$. Then in both 7.8.2 and 7.8.3, Γ_1 has $\Lambda_0 \times \Lambda_0$ as a relatively rigid subgroup, which is wq-normal in Γ_1 , with $\Lambda \times \Lambda$ ICC in Γ_1 . Furthermore, since $\text{Out}(\mathcal{R}_{\sigma_i})$ is huge for $i \geq 2$ and trivial for $i = 1$, σ_1 cannot be stably orbit equivalent to $\sigma_i, i \geq 2$. The existence of “many actions” σ in the case 7.8.2 and 7.8.3 follows from the existence of uncountably many non-stably OE relations (σ_1, Γ_1) of the form 7.8.2 (cf [MS]) and respectively of the form 7.8.3 (by Proposition 7.9.3° and [MS]).

Finally, the calculation of the 1-cohomology groups follows from (2.12, 3.1, 3.2 in [P8]) and the fact that in all cases 7.8.1, 7.8.2, 7.8.3 one has $H^1(\sigma_1, \Gamma_1) = \text{Char}(\Gamma_1)$. Indeed, in case σ_1 is as in 7.8.1 or 7.8.2 then this calculation follows from ([Ge1] and 2.12 in [P8]), while in case 7.8.3 the calculation is in ([P8]). $\square$

7.11. Corollary. *With $\Gamma = *_{i \geq 0} \Gamma_i$, K , σ as in 7.10.2°, denote $A = L^\infty(X, \mu)$ and $M = A \rtimes_\sigma \Gamma$. Then $\mathcal{F}(M) = \{1\}$ and*

$$\text{Out}(M) = H^1(\sigma, \Gamma) = \mathbb{G}_0^{n-1} \times \mathbb{G} \times \prod_{j \geq 1} \text{Char}(\Gamma_j) \times K^{n-1}.$$

Proof. Trivial by 7.7 and 7.10. $\square$

Note that $\text{Out}(M)$ is abelian and non-locally compact in all examples in 7.11 above, but if we denote $\tilde{\text{Out}}(M)$ the quotient of $\text{Out}(M)$ by the connected component of id_M (which is closed in $\text{Out}(M)$, with the latter a polish group in all examples considered), then $\tilde{\text{Out}}(M)$ is the quotient of $\prod_{j \geq 1} \text{Char}(\Gamma_j) \times K^{n-1}$ by the connected component of 1, which for $n < \infty$ is totally disconnected separable, locally compact group.

We end this section by mentioning another rigidity result, which from an isomorphism of group measure space factors corresponding to relatively rigid actions ([P5]) of free products of groups derives the orbit equivalence of the actions. This type of results were first obtained in ([P5]) for HT group-actions, and in ([P6]), for Bernoulli shift actions of groups containing infinite subgroups (not necessarily normal) with the relative property (T).

7.12. Theorem (vNE/OE rigidity). *Let (M_i, τ_i) be type II_1 von Neumann algebras with a common Cartan subalgebra $A \subset M_i$, $i = 1, 2$, such that $\tau_1|_A = \tau_2|_A$. Assume $M = M_1 *_A M_2$ is a factor and A is Cartan in M (N.B. By 7.2 this is the same as requiring that $\mathcal{R}_{A \subset M_i}$, $i = 1, 2$, are freely independent). If $A_0 \subset M^t$ is a rigid Cartan subalgebra, for some $t > 0$, then there exists a unitary element $u \in M^t$ such that $uA_0u^* = A^t$.*

Proof. It is clearly sufficient to prove this in the case $t = 1$. If some corner of A_0 can be embedded into A inside M , then the statement follows by (A.1 in [P5]). If we assume this is not the case, then we can apply Theorem 7.5 to get a non-zero $p \in \mathcal{P}(A_0)$ such that $vA_0pv^* \subset M_i$ for some $i \in \{1, 2\}$ and $v \in \mathcal{U}(M)$. Moreover, since M is a factor and A_0 is Cartan in M , we may assume vpv^* is central in M_i , so in particular $p_1 = vpv^*$ lies in A (the latter being maximal abelian in M_i). Then vA_0pv^* is Cartan in p_1Mp_1 , so by Corollary 1.2.4 we have $p_1M_2p_1 = Ap_1$, contradicting the fact that M_2 is type II . $\square$

7.13. Corollary. *Let $\mathcal{F}$ be the class of groups that can be written as a free product of two (or more) infinite groups. Let $\sigma : G \rightarrow (X, \mu)$ be a free ergodic m.p. action of a group $G \in \mathcal{F}$ and denote $M = L^\infty(X, \mu) \rtimes_\sigma G$ the corresponding group measure space II_1 factor, with $A = L^\infty(X, \mu) \subset M$ the corresponding Cartan subalgebra. Let M_0 be a II_1 factor with a relatively rigid Cartan subalgebra $A_0 \subset M_0$ and denote $\mathcal{R}_0 = \mathcal{R}_{A_0 \subset M_0}$.*

1°. *If $\theta : M_0 \simeq M^t$ for some $t > 0$ then θ can be perturbed by an inner automorphism so that to take A_0 onto A^t . In particular, $\mathcal{R}_0 \simeq \mathcal{R}_\sigma^t$ and thus $\beta_n(\mathcal{R}_0) = \beta_n(G)/t$, $\forall n$.*

2°. *If $G \in \mathcal{F}$ satisfies $\beta_n(G) \neq 0, \infty$ for some n (for instance, if $G = \Gamma_1 * \Gamma_2$, with $\Gamma_{1,2}$ finitely generated infinite groups, in which case $\beta_1(G) \neq 0, \infty$) and the action σ is relatively rigid, then $\mathcal{F}(M) = \{1\}$.*

Proof. All statements are trivial by 7.12 and ([G1]). $\square$

Note that 7.13.2° above shows in particular that $\mathcal{F}(L(\mathbb{Z}^2 \rtimes \mathbb{F}_n)) = \{1\}$ for any $\mathbb{F}_n \subset SL(2, \mathbb{Z})$, with $2 \leq n < \infty$, thus giving a new proof (but still using Gaboriau's work [G1]) of one of the main result in ([P5]).

7.14. Definition. A countable, measurable, standard m.p. equivalence relations $\mathcal{R}$ is a *FT equivalence relation* if it is of the form $\mathcal{R} = \mathcal{R}_\sigma^t$, where $t > 0$ and (σ, Γ) are free ergodic m.p. actions on the probability space (X, μ) with the following properties: (a). The group Γ is a free product of two (or more) infinite groups; (b). σ is relatively rigid, in the sense of (Definition 5.10.1 in [P5]), i.e. $L^\infty(X, \mu) \subset L^\infty(X, \mu) \rtimes_\sigma G$ is a rigid inclusion (4.2.1 in [P5]). The above Corollary 7.13 thus shows that all OE invariants for *FT* equivalence relations $\mathcal{R}$ are in fact vNE invariants for $\mathcal{R}$, i.e. are isomorphism invariants of the associated group measure space II_1 factors $M = L(\mathcal{R}, w)$, where $w \in H^2(\mathcal{R})$. We denote by $\mathcal{FT}$ the class of all such II_1 factors M .

Note that if $\mathcal{R} = \mathcal{R}_{\sigma, \Gamma}^t$ for some free ergodic m.p. action (σ, Γ) with Γ a free product of two infinite groups, then $\mathcal{R}$ is *HT_s* in the sense of ([P5]) if and only if it is *FT* and Γ has Haagerup property. Thus, all equivalence relations coming from amplifications of actions σ of non-amenable subgroups $\Gamma \subset SL(2, \mathbb{Z})$ on $L^\infty(\mathbb{T}^2, \lambda)$ are *FT* actions. However, actions σ of groups such as $SL(n, \mathbb{Z}) * H$, with H infinite group and $\sigma|_{SL(n, \mathbb{Z})}$ isomorphic

to the canonical action of $SL(n, \mathbb{Z})$ on $(\mathbb{T}^n, \lambda)$, give FT equivalence relations which are not HT_s . Thus, the class $\mathcal{FT}$ provides additional group measure space II_1 factors for which orbit equivalence invariants of the actions, such as Gaboriau's ℓ^2 -Betti numbers, become isomorphism invariants of the factors.

We end by mentioning an application of 7.3.1° which brings some light to (Problems 5.10.2, 6.12.1 in [P5]) and to the problem of existence of “many” non-OE actions for non-amenable groups, as a consequence of (Corollary 7 in [GP]):

7.15. Corollary. 1°. *The class of groups Γ which admit free ergodic relatively rigid m.p. actions on the probability space is closed to free products with arbitrary groups Γ' . Also, if Γ is a H_{T_s} group (i.e. Γ has Haagerup's property and admits relatively rigid actions, see 6.11 in [P5]), then $\Gamma * \Gamma'$ is H_{T_s} , for any Γ' with Haagerup property.*

2°. *If Γ admits a relatively rigid action (e.g. if $\Gamma \subset SL(2, \mathbb{Z})$ non-amenable, or Γ an arithmetic lattice in an absolutely simple, non-compact Lie group with trivial center, cf [P5], [Va]) and Γ' is an arbitrary infinite amenable group then $\Gamma * \Gamma'$ has uncountably many non-stably OE free ergodic m.p. actions on the probability space. Also, if Γ_0 is an arbitrary group and Γ_1, Γ_2 are non-trivial amenable groups, at least one having more than 2 elements, then $\Gamma_0 * \Gamma_1 * \Gamma_2$ has uncountably many non-stably OE free ergodic m.p. actions on the probability space.*

Proof. Part 1° is a trivial consequence of 7.3.1° and of the (trivial) property of relatively rigid equivalence relations $\mathcal{R}$ that any $\mathcal{R}_0$ that contains $\mathcal{R}$ is also relatively rigid (see e.g. 4.6.2° in [P5]). Part 2° is just the combination of (Corollary 7 in [GP]) and 7.3. $\square$

8. AMALGAMATION OVER R : FACTORS WITH NO OUTER AUTOMORPHISMS

We prove here another rigidity result for AFP factors, this time in the case $M = M_0 *_R M_1 *_R \dots$, where R is the hyperfinite II_1 factor. As an application, we obtain factors M with $\text{Out}(M) = \{1\}$, thus answering a well known problem posed by A. Connes in 1973.

Like in the group measure space case in Section 7, we only consider crossed product inclusions $(R \subset M_i) = (R \subset R \rtimes_{\sigma_i} \Gamma_i)$, with the σ_i freely independent, i.e. implementing a free action σ of $\Gamma = \Gamma_0 * \Gamma_1 * \dots$ on R . Thus, M will be viewed alternatively as a crossed product factor $M = R \rtimes_{\sigma} \Gamma$, with the algebra of coefficients R having trivial relative commutant in M .

The key assumption is that the action $(\sigma, *_i \Gamma_i)$ has the relative property (T), i.e. $R \subset M$ be a rigid inclusion in the sense of ([P5]). The rigidity result shows the uniqueness, modulo unitary conjugacy, of the “core” R of such factors. Since the normalizer of R in M completely encodes the group Γ , we can recover completely the isomorphism class of the groups Γ_i , by classical Bass-Serre theory. The result is similar to the vNE/OE Rigidity 7.11 (where however only the orbit equivalence class of Γ could be recovered) and to the unique crossed-product decomposition result in ([P9]). But since we also get the component by component unitary conjugacy of the factors M_i , it is again a Bass-Serre type rigidity result.

Through this theorem, the calculation of $\text{Out}(M)$ and $\mathcal{F}(M)$ reduces to the calculation of the commutant of $\sigma(\Gamma)$ in $\text{Out}(R)$, like in [P9], where however no such commutant could

be calculated! This time, due to Bass-Serre arguments and the possibility of choosing the actions (σ_i, Γ_i) with prescribed properties (cf. 8.2 below), we can control such commutants and calculate $\text{Out}(M)$ completely for large classes of factors.

8.1. Lemma. *Let $M_n, n \geq 0$, be II_1 factors with a common subfactor $N \subset M_n$. Then $N \subset M = M_0 *_N M_1 *_N M_2 *_N \dots$ is irreducible and regular if and only if $N \subset M_n$ is regular, irreducible $\forall n \geq 1$ and the groups of outer automorphisms $\Gamma_n = \{\text{Ad}(u) \mid u \in \mathcal{N}_{M_n}(N)\}/\mathcal{U}(N)$, $n \geq 0$, on N are freely independent.*

Proof. Trivial by the definitions of freeness and respectively of the amalgamated free product over N . $\square$

The result below is the analogue for actions on the hyperfinite II_1 factor R of the result on the existence of freely independent actions on the probability space in Proposition 7.3. It shows the existence of free actions σ of groups $\Gamma = \Gamma_0 * \Gamma_1 * \Gamma_2 * \dots$ on R such that the restriction of σ to each individual group Γ_j is conjugate to a prescribed free action of Γ_j on R . It will be frequently used in this Section. The proof relies on Lemma A.2 in the Appendix.

8.2. Proposition. *Let $\sigma_n : \Gamma_n \rightarrow \text{Aut}(R)$ be free actions of countable discrete groups G_n , $n \geq 0$. Then there exists a free action σ of the group $G = *_n \Gamma_n$ on R such that $\sigma|_{\Gamma_n}$ is conjugate to σ_n , $\forall n \geq 0$.*

Proof. For each $n \geq 0$ denote $\tilde{G}_n = G_0 * G_1 * \dots * G_n$. Assume we have constructed a map $\tilde{\sigma}_n$ of $\tilde{G}_n$ into $\text{Aut}(R)$ such that the quotient map $\tilde{\sigma}'_n$ of G into $\text{Aut}(R)/\text{Int}(R)$ is a faithful group morphism with $\tilde{\sigma}_n|_{G_i}$ conjugate to σ_i , $\forall 0 \leq i \leq n$. We then apply A.2 to $\{\tilde{\sigma}_n(g) \mid g \in \tilde{G}_n\} \cup \{\sigma_{n+1}(h) \mid h \in G_{n+1}\}$ to get an automorphism θ_{n+1} of R such that $\tilde{\sigma}_n(\tilde{G}_n)$ and $\theta_{n+1}\sigma_{n+1}(G_{n+1})\theta_{n+1}^{-1}$ are freely independent. Denoting by $\tilde{\sigma}_{n+1}$ the map of $\tilde{G}_{n+1} = \tilde{G}_n * G_{n+1}$ into $\text{Aut}(R)$ which restricted to $\tilde{G}_n$ equals $\tilde{\sigma}_n$ and restricted to G_{n+1} equals $\theta_{n+1}\sigma_{n+1}(G_{n+1})\theta_{n+1}^{-1}$, the statement follows by induction. $\square$

8.3. Theorem. *Let $G_i = \Gamma_{i,0} * \Gamma_{i,1} * \dots * \Gamma_{i,n_i}$ with $\Gamma_{ij}, 0 \leq j \leq n_i$, non-trivial groups, for some $1 \leq n_i \leq \infty$, $i = 1, 2$. For each $i = 1, 2$ let $\sigma_i : G_i \rightarrow \text{Aut}(N_i)$ be a free, ergodic m.p. action on a II_1 factor N_i . Denote $M_i = N_i \rtimes_{\sigma_i} G_i$, $i = 1, 2$ and assume $N_i \subset M_i$ are rigid inclusions $i = 1, 2$. Let $\theta : M_1 \simeq M_2^t$, for some $t > 0$. Then we have:*

1°. *There exists $u \in \mathcal{U}(M_2^t)$ such that $\text{Ad}(u)(\theta(N_1)) = N_2^t$. Thus, $G_1 \simeq G_2$ and σ_1, σ_2^t are cocycle conjugate actions with respect to the identification $G_1 \simeq G_2$.*

2°. *If in addition $\Gamma_{i,0}$ are free groups and Γ_{ij} are free indecomposable, not equal to the infinite cyclic group, $\forall 1 \leq j \leq n_i$, $i = 1, 2$, then $\Gamma_{10} \simeq \Gamma_{20}$, $n_1 = n_2$ and there exists a permutation π of the indices $j \geq 1$ and unitaries $u_j \in M_2^t$ such that $\text{Ad}(u_j)(\theta(M_{1,j})) = M_{2,\pi(j)}^t$, $\text{Ad}(u_j)(\theta(N_1)) = N_2^t$, $\forall j \geq 1$. In particular, $\Gamma_{1,j} \simeq \Gamma_{2,\pi(j)}$ and $\sigma_{1,j}, \sigma_{2,\pi(j)}^t$ are cocycle conjugate with respect to this identification of groups, $\forall j \geq 1$.*

Proof. We first prove that a corner of $\theta^{-1}(N_2^t)$ can be embedded into N_1 inside M_1 . Assume this is not the case. By applying recursively 5.1.3°, it follows that there exist a unitary

element $u \in \mathcal{U}(M_1)$ and some $n_1 \geq j \geq 1$ such that $u\theta^{-1}(N_2^t)u^* \subset M_{1,j}$. Since $u\theta^{-1}(N_2^t)u^*$ is regular in M_1 , using again the assumption by contradiction, Corollary 1.2.4 implies that a corner of $u\theta^{-1}(N_2^t)u^*$ can be embedded into N_1 inside $M_{1,j}$ (thus inside M_1 as well). Altogether, this shows that a corner of $\theta^{-1}(N_2^t)$ can be embedded into N_1 inside M_1 .

Similarly, a corner of $\theta(N_1)^{1/t} = \theta^{1/t}(N_1^{1/t})$ can be embedded into N_2 inside M_2 . Thus, a corner of N_1 can be embedded into $\theta^{-1}(N_2^t)$ inside M_1 . Since both N_1 and $\theta^{-1}(N_2^t)$ are regular in M_1 , with $\mathcal{N}_{M_1}(N_1)/\mathcal{U}(N_1) \simeq G_1$ and with the other similar quotient isomorphic to G_2 , and since both G_1, G_2 are ICC (being free products of non-trivial groups), the unitary conjugacy of $N_1, \theta^{-1}(N_2^t)$ in M_1 (equivalently of $\theta(N_1), N_2^t$ in M_2^t) follows from the following general:

8.4. Lemma. *Let M be a II_1 factor and $P, Q \subset M$ be irreducible, regular subfactors. Assume $\Gamma = \mathcal{N}_M(P)/\mathcal{U}(P)$, $\Lambda = \mathcal{N}_M(Q)/\mathcal{U}(Q)$ are ICC groups. Also, assume each one of the inclusions $P \subset M$, $Q \subset M$ is an amplification of a genuine crossed product inclusion. If $L^2(M)$ contains non-zero $P - Q$ Hilbert bimodules $\mathcal{H}, \mathcal{K} \subset L^2(M)$ such that $\dim_P(\mathcal{H}) < \infty$, $\dim(\mathcal{K})_Q < \infty$ then P, Q are unitary conjugate in M .*

Proof. We first prove that $L^2(M)$ is generated by irreducible $P - Q$ Hilbert bimodules that are finite dimensional both as left P modules and as right Q modules. We'll actually prove this by only using the fact that P, Q are quasi-regular in M . Note that by (1.4 in [P5]) $\mathcal{H}^0 = \mathcal{H} \cap M$ is dense in $\mathcal{H}$ and contains an orthonormal basis/ Q . Similarly, since Q is quasi-regular in M and it is a factor, $L^2(M)$ is generated by Hilbert $Q - Q$ bimodules $\mathcal{H}_\beta$ such that $\mathcal{H}_\beta^0 = \mathcal{H}_\beta \cap M$ is dense in $\mathcal{H}_\beta$ and contains both left and right orthonormal basis/ Q . But then $\mathcal{H}^0 \cdot \mathcal{H}_\beta^0$ span all $L^2(M)$ and are finite dimensional/ Q . Equivalently, $P' \cap J_M Q' J_M$ is generated by projections that have finite trace in $J_M Q' J_M$. Similarly, $P' \cap J Q' J$ is generated by projections that have finite trace in P' . Thus, $\mathcal{A} = P' \cap J Q' J$ is generated by projections that are finite with respect to both traces, thus corresponding to Hilbert $P - Q$ bimodules which are finite dimensional both from right and left. Since P, Q are factors, by ([J]) each such bimodule is a direct sum of irreducible bimodules.

Let now $\mathcal{H} \subset L^2(M)$ be an irreducible $P - Q$ bimodule. By ([J]) we have $\dim({}_P \mathcal{H}) \cdot \dim(\mathcal{H}_Q) \geq 1$ and the equality means the orthogonal projection $p_{\mathcal{H}}$ of $L^2(M)$ onto $\mathcal{H}$ satisfies: $p_{\mathcal{H}} \in P' \cap \langle M, e_Q \rangle$, $\text{Tr}_{\langle M, Q \rangle}(p_{\mathcal{H}}) = 1$ and $p_{\mathcal{H}} \langle M, e_Q \rangle p_{\mathcal{H}} = P p_{\mathcal{H}}$. Thus, by the proof of (Lemma 1 in [P7]) $u p_{\mathcal{H}} u^* = e_Q$ for some $u \in \mathcal{U}(M)$, which also satisfies $u P u^* = Q$.

Assume now that $\text{Tr}(p_{\mathcal{H}}) > 1$, by (2.1 in [P6]) there exists a projection $p \in P$, a unital isomorphism $\psi : p P p \rightarrow Q$ and a partial isometry $v \in M$ such that $vv^* = p$, $q' = v^* v \in \psi(p P p)' \cap Q$ and $xv = v\psi(x)$, $\forall x \in p P p$. Moreover, the finite dimensionality plus irreducibility of $\mathcal{H}$ as a $P - Q$ bimodule implies that $Q_1 = \psi(p P p)$ has finite index in Q , has trivial relative commutant in Q and q' is minimal in $Q'_1 \cap M$.

By appropriately amplifying $Q \subset M$ we may assume this inclusion is a genuine crossed product inclusion $Q \subset Q \rtimes_{\sigma} \Gamma$. Denote by $\{u_g\}_g \subset M = Q \rtimes_{\sigma} \Gamma$ the canonical unitaries implementing σ on Q . Let $q' = \sum_g x_g u_g$, with $x_g \in Q$. By identification of Fourier series, it follows that $x_g u_g \in Q'_1 \cap M$, $\forall g$. Thus $x_g x_g^* = x_g u_g u_g^* x_g^* \in Q'_1 \cap Q = \mathbb{C}$, so that all x_g are scalar multiples of unitaries in Q . Let $K_0 \subset \Gamma$ be the support of this Fourier expansion of

q' . Let also $K \subset \Gamma$ be the set of all $k \in \Gamma$ such that u_k can be perturbed a unitary in Q so that to fix Q_1 pointwise. Since $[Q : Q_1] < \infty$ and $Q'_1 \cap Q = \mathbb{C}$, K is a finite subgroup of Γ and $K_0 \subset K$.

Since $Q'_1 \cap Q = \mathbb{C}$, by Connes' vanishing 1-cocycle for finite groups, the unitaries $w_k \in Q$ satisfying $\text{Ad}(w_k u_k)|_{Q_1} = \text{id}_{Q_1}$ can be chosen of the form $\sigma_k(w)w^*$, $k \in K$, for some unitary element $w \in Q_1$. Thus, by perturbing all $\{u_g\}_{g \in \Gamma}$ by a 1-cocycle we may assume $\text{Ad}(u_k)$ act trivially on Q_1 . Let $\Gamma_0 \subset \Gamma$ be the subgroup of all $g \in \Gamma$ such that u_g can be perturbed by a unitary in Q so that to normalize Q_1 . Clearly $K \subset \Gamma_0$ and K is normal in Γ_0 . We will prove that Γ_0 has finite index in Γ , thus contradicting the hypothesis.

By the minimality of q' in $Q'_1 \cap M$, it follows that q' is minimal in the group algebra $L(K) = \text{sp}\{u_k \mid k \in K\}$. Identify $pPp \subset pMp$ with $Q_1 q' \subset q' M q'$ via $\text{Ad}(v)$. Let $\{v_h \mid h \in \Lambda\}$ be a choice of canonical unitaries in $M = P \rtimes \Lambda$, which we assume commute with $p \in P$ (we can do that for each h by perturbing if necessary with unitaries in the factor P). For each $h \in \Lambda$, $h \neq e$, let $v_h = \sum_g x_g^h u_g \in Q \rtimes_\sigma \Gamma$ be the Fourier expansion of v_h and denote θ_h the action implemented by $\text{Ad}(v_h)$ on $Q_1 \simeq Q_1 q' = pPp$. Thus, $v_h y = \theta_h(y) v_h, \forall y \in Q_1$. Identifying the Fourier series in $\{u_g\}_g$, this implies $\theta_h(y)(x_g^h u_g) = (x_g^h u_g) y, \forall y \in Q_1$. As before, this implies each x_g^h is a scalar multiple of a unitary in Q and that $\text{Ad}(x_g^h u_g)$ normalizes Q_1 . Thus the support K_h of the Fourier series for v_h is contained in Γ_0 . Since $Q_1 q' \subset q' M q'$ is the closure of the span of elements in $Q_1 v_h q', h \in \Lambda$ and each v_h is supported on Γ_0 , as a Fourier expansion in $\{u_g\}_g$ with coefficients in Q , it follows that $q' M q' \subset \bigoplus_{g \in \Gamma_0} L^2(Q) u_g$. In particular, since $q' \in L(K) \subset L(\Gamma_0)$ we get $q' L(\Gamma) q' = q' L(\Gamma_0) q'$, which clearly implies Γ_0 has finite index in Γ . But this implies Γ_0 is also ICC, so in particular it cannot have a non-trivial normal subgroup K . This contradiction finishes the proof. $\square$

End of proof of 8.3. By Lemma 8.4, $\theta(N_1), N_2^t$ are conjugate by a unitary element, so we may assume $\theta(N_1) = N_2^t$. Thus, θ implements an isomorphism between the groups $\Gamma_1 = \mathcal{N}_{M_1}(N_1)/\mathcal{U}(N_1)$ and $\Gamma_2 = \mathcal{N}_{M_2}(N_2)/\mathcal{U}(N_2)$. But then by the classical Kurosh Theorem and the condition on “free indecomposability” of the groups Γ_{ij} , it follows that $n_1 = n_2 = n$ and that there exists a permutation π of the indices $1 \leq j \leq n$ such that $g_j \Gamma_{1,j} g_j^{-1} = \Gamma_{2,\pi(j)}$, for some elements $g_i \in G$. Thus, $u_j = u_{g_j}$ normalizes N_2^t and $\text{Ad}(u_j)$ takes $\theta(M_{1,j})$ onto $M_{2,\pi(j)}^t$. $\square$

8.5. Notation. We denote by $f\mathcal{T}_R$ the class of free actions $\sigma : \Gamma_0 * \Gamma_1 \rightarrow \text{Aut}(R)$ on the hyperfinite II_1 factor R , with the properties:

(8.5.1). Γ_0 is free indecomposable; Γ_1 is w-rigid (in particular free indecomposable).

(8.5.2). $\sigma_0 = \sigma|_{\Gamma_0}$ has the relative property (T), i.e. $R \subset R \rtimes_{\sigma_0} \Gamma_0$ is a rigid inclusion; $\sigma_1 = \sigma|_{\Gamma_1}$ is a non-commutative Bernoulli shift action of Γ_1 on $R = \overline{\otimes}_g (N_0, \tau_0)_g$, where $N_0 = R$ or $N_0 = M_{n \times n}(\mathbb{C})$ for some $n \geq 2$.

(8.5.3). $\sigma(\Gamma_1)$ and the normalizer of $\sigma(\Gamma_0)$ in $\text{Out}(R)$ (which is countable by [P5]) are free independent.

8.6. Lemma. *Let Γ_1 be an arbitrary w -rigid group and $\Gamma_0 = SL(n, \mathbb{Z})$, $n \geq 2$, more generally Γ_0 a free indecomposable, arithmetic lattice in an absolutely simple, non-compact Lie group with trivial center. Then $\Gamma_0 * \Gamma_1$ has $f\mathcal{T}_R$ actions on R .*

Proof. By ([Va]) any such Γ_0 has a free ergodic action on some $\mathbb{Z}^m$ such that the pair $(\mathbb{Z}^m \rtimes \Gamma_0, \mathbb{Z}^m)$ has the relative property (T) of Kazhdan-Margulis ([Ma]). By ([Ch], [NPS]) it follows that Γ_0 admits a free action σ_0 on the hyperfinite II_1 factor R such that $R \subset R \rtimes_{\sigma_0} \Gamma_0$ has the relative property (T). By ([P5]) it follows that the normalizer $\mathcal{N}_0$ of $\sigma_0(\Gamma_0)$ in $\text{Out}(R)$ is countable. Let σ' be a fixed copy Bernoulli shift action of Γ_1 . By 8.2 it follows that there exists an automorphisms θ of R such that $\theta(\sigma'(\Gamma_1))\theta^{-1}$ is freely independent from $\mathcal{N}_0$.

Thus, if we denote by σ the unique action of $\Gamma = \Gamma_0 * \Gamma_1$ on R given by $\sigma|_{\Gamma_0} = \sigma_0$ and $\sigma|_{\Gamma_1} = \theta\sigma'\theta^{-1}$, then all conditions (8.5.1) – (8.5.3) are satisfied. $\square$

8.7. Theorem. *Let $\sigma_i : \Gamma_{i0} * \Gamma_{i1} \rightarrow \text{Aut}(R_i)$, $i = 1, 2$, be $f\mathcal{T}_R$ actions and denote $G_i = \Gamma_{i0} * \Gamma_{i1}$, $M_i = R_i \rtimes_{\sigma_i} G_i$, $i = 1, 2$. If $\theta : M_1 \simeq M_2^t$ is an isomorphism, for some projection $t > 0$ then $t = 1$ and there exist a unitary $u \in M_2$, a character γ of Γ_{20} and isomorphisms $\delta : G_1 \simeq G_2$, $\Delta : R_1 \simeq R_2$ such that $\theta_0 = \text{Ad}(u) \circ \theta^\gamma \circ \theta$ satisfies. Moreover, any other isomorphism $\theta' : M_1 \simeq M_2$ is a perturbation of θ by an automorphism of M_2 of the form $\text{Ad}(v) \circ \theta^{\gamma'}$ for some $v \in \mathcal{U}(M_2)$ and $\gamma' \in \text{Char}(G_2)$.*

Proof. Since $R_i \subset R_i \rtimes_{\sigma_i} \Gamma_{i0}$ are rigid inclusions, $R_i \subset R_i \rtimes_{\sigma_i} G_i = M_i$ are rigid as well. We can thus apply Theorem 8.3 to get a unitary element $v \in M_2^t$ such that $v(\theta(R_1))v^* = R_2^t$ and $v(\theta(M_{11}))v^* = (M_{2j})^t$ for some $j \in \{0, 1\}$, where $M_{ij} = R_i \rtimes \Gamma_{ij}$, $i = 1, 2, j = 0, 1$. But by ([P5] or [P6]) $R_1 \subset M_{11}$ is not rigid while $(R_2 \subset M_{20})^t$ is rigid, so the only possibility is that $j = 1$. Thus, $(R_1 \subset M_{11}) \simeq (R_2 \subset M_{21})^t$ and both inclusions come from crossed products associated with non-commutative Bernoulli shift actions of w -rigid groups. By ([P3]) this implies $t = 1$.

On the other hand, $\text{Ad}v \circ \theta$ implements an isomorphism $\delta : \Gamma_{10} * \Gamma_{11} \simeq \Gamma_{20} * \Gamma_{21}$, which takes Γ_{11} onto Γ_{21} . By Kurosh Theorem, $\delta(\Gamma_{10}) = g\Gamma_{20}g^{-1}$ for some $g \in G_2$. But by ([GoS]), the groups $g\Gamma_{20}g^{-1}$ and Γ_{21} can generate $\Gamma_{20} * \Gamma_{21}$ only if $g = g_1g_2$ for some $g_i \in \Gamma_{2i}$, $i = 0, 1$. By conjugating with g , we may thus assume the unitary v is so that $\theta_0 = \text{Ad}(v) \circ \theta$ implements an isomorphism $\delta : G_1 \simeq G_2$ that takes Γ_{1j} onto Γ_{2j} , $j = 0, 1$. Thus, after identifying R_1 with R_2 via $\Delta = (\theta_0)|_{R_1}$ and $G_1 \simeq G_2$ via δ , we are left with finding all automorphisms α of M_2 that take R_2 onto itself and take the canonical unitaries u_g into unitaries $w_g u_g$, $g \in G_2$, for some $w : G_2 \rightarrow \mathcal{U}(R_2)$ a 1-cocycle for σ_2 .

By ([P3]) this implies w is co-boundary when restricted to Γ_{21} , i.e. $w_g = \sigma_g(w)w^*$, $\forall g \in \Gamma_{21}$, for some unitary $w \in R_2$. Thus, by replacing α by $\text{Ad}(v^*) \circ \alpha$, we may assume $\alpha(u_g) = u_g, \forall g \in \Gamma_{21}$. Thus $\alpha|_{R_2} \in \text{Aut}(R_2)$ commutes with $\sigma_2(\Gamma_{21})$, while still normalizing $\sigma_2(\Gamma_{20})$. But by condition (8.5.3) the latter condition implies $\alpha|_{R_2}$ is freely independent from $\sigma_2(\Gamma_{21})$. This contradicts the commutation condition with $\sigma_2(\Gamma_{21})$, unless $\alpha|_{R_2}$ is inner. By perturbing α by $\text{Ad}(w_0)$ for an appropriate $w_0 \in \mathcal{U}(R_2)$ we may thus assume $\alpha_{R_2} = \text{id}_{R_2}$. Thus α is given by a character of G_2 . $\square$

The above theorem shows in particular that the fundamental group of any $f\mathcal{T}_R$ factor

$M = R \rtimes_{\sigma} (\Gamma_0 * \Gamma_1)$, corresponding to an $f\mathcal{T}_R$ action $(\sigma, \Gamma_0 * \Gamma_1)$, is trivial, while its Out-group is equal to $\text{Char}(\Gamma_0) \times \text{Char}(\Gamma_1)$. By Lemma 8.6 one can take $\Gamma_0 = SL(n, \mathbb{Z})$, which has only trivial characters, thus making $\text{Aut}(M) = \text{Char}(\Gamma_1)$, with Γ_1 an arbitrary w-rigid group. For instance, one can take $\Gamma_1 = SL(3, \mathbb{Z}) \times H$, where H is an arbitrary discrete abelian group, thus getting $\text{Aut}(M) = \hat{H}$. We thus obtain:

8.8. Corollary. *Let $\sigma : \Gamma_0 * \Gamma_1 \rightarrow \text{Aut}(R)$ be a $f\mathcal{T}_R$ action and denote $M = R \rtimes_{\sigma} (\Gamma_0 * \Gamma_1)$. Then we have:*

- 1°. $\mathcal{F}(M) = \{1\}$ and $\text{Out}(M) = \text{Char}(\Gamma_0) \times \text{Char}(\Gamma_1) = \text{Out}(M^{\infty})$.
- 2°. *Given any compact abelian group K , there exists $(\sigma, \Gamma_0 * \Gamma_1)$ such that the corresponding $f\mathcal{T}_R$ factor M satisfies $\text{Out}(M) = K = \text{Out}(M^{\infty})$. For instance, if $\Gamma_0 = SL(n, \mathbb{Z})$, $\Gamma_1 = SL(m, \mathbb{Z}) \times \hat{K}$ for some $n, m \geq 3$, then $\text{Out}(M) = K$.*

8.9. Remark. One can use A.3.2° in lieu of A.2 in all the above proofs, to construct more II_1 factors with small, calculable symmetry groups. Thus, let $f\mathcal{T}'_R$ be the class of free actions $\sigma : \Gamma_0 * \Gamma_1 \rightarrow \text{Aut}(R)$ on the hyperfinite II_1 factor R , satisfying the properties:

- (a). Γ_0, Γ_1 free indecomposable, $\neq \mathbb{Z}$;
- (b). $R \subset R \rtimes_{\sigma_0} \Gamma_0$ is a rigid inclusion and $\sigma_1 = \sigma|_{\Gamma_1}$ is non-cocycle conjugate to $\sigma_0 = \sigma|_{\Gamma_0}$ (note that this is indeed the case if σ_1 is a non-commutative Bernoulli shift);
- (c). $\sigma(\Gamma_1)$ is free independent with respect to the set $\mathcal{N}(\sigma_0(\Gamma_0)) \cup \mathcal{N}^{op}(\sigma_0(\Gamma_0))$, consisting of all automorphisms and anti-automorphisms of R^{∞} that normalize $\sigma_0(\Gamma_0)$.

By A.3.2° and 8.6 it follows that there exist such actions σ for any linear group Γ_0 as in 8.6 and for any Γ_1 free indecomposable. It then follows as in 8.8 that the corresponding crossed product II_1 factors $M = R \rtimes_{\sigma} (\Gamma_0 * \Gamma_1)$ satisfy $\text{Out}(M) = \text{Char}(\Gamma_0 * \Gamma_1)$. Moreover, if $t \in \mathcal{F}(M)$ and $\theta : M \simeq M^t$ then θ must normalize $\sigma(\Gamma_0)$, contradicting condition (c) above. Thus, $\mathcal{F}(M) = \{1\}$. Notice that to get this calculation we no longer have to use the results in ([P3]) on the fundamental group of $R \subset R \rtimes_{\sigma_1} \Gamma_1$. Similarly, if α is an anti-automorphism of M then the same argument shows that it must normalize $\sigma_0(\Gamma_0)$, in contradiction with the choice (c). Altogether, this shows that in addition to the properties $\mathcal{F}(M) = \{1\}$, $\text{Out}(M) = \text{Char}(\Gamma_0 * \Gamma_1)$, the factors M in the class $f\mathcal{T}'_R$ have no anti-automorphisms either. This provides a fairly large new family of factors with this latter property, after Connes first examples in ([C4]). Thus, if we choose the groups Γ_0, Γ_1 without characters, e.g. $\Gamma_i = SL(n_i, \mathbb{Z})$, $n_i \geq 3$, then the resulting factors M have no outer symmetries at all. Moreover, noticing that $f\mathcal{T}'_R$ factors are w-rigid, it follows that any factor of the form $N * M$, with N a property (T) II_1 factor (e.g. $N = L(PSL(n, \mathbb{Z}))$, $n \geq 3$) and $M \in f\mathcal{T}'_R$ has all the above properties (as a consequence of Theorem 6.3 and of the properties of M) and in addition has no Cartan subalgebras (by [V2], cf. Remark 6.6)!

APPENDIX: CONSTRUCTING FREELY INDEPENDENT ACTIONS.

We now prove the technical results needed in the proofs of Propositions 7.3 and 8.2, which established the existence of free actions of groups $\Gamma = \Gamma_0 * \Gamma_1 * \dots$ on $A = L^{\infty}(X, \mu)$ and on R with restrictions to Γ_i isomorphic to prescribed actions (σ_i, Γ_i) , $\forall i$.

More precisely, we prove that given any countable set $\{\theta_n\}_n$ of properly outer automorphisms of (X, μ) (resp. of R) the set $\mathcal{V}$ of $\theta \in \text{Aut}(X, \mu)$ (resp $\theta \in \text{Aut}(R)$) with the property that all alternating words $\theta_{i_0} \theta \theta_{i_1} \theta^{-1} \theta_{i_2} \theta \theta_{i_3} \theta^{-1} \dots$ are properly outer is G_δ -dense in $\text{Aut}(X, \mu)$ (resp. $\text{Aut}(R)$). Writing $\mathcal{V}$ as an intersection of open sets $\mathcal{V}_n$ is obvious, and the non-trivial part is to show that each $\mathcal{V}_n$ is dense. To prove the density, in the commutative case (A.1) we use a maximality argument inspired from ([P7]), while in the hyperfinite case (A.2) we use directly a result from ([P7]), not having to re-do such maximality argument.

The idea of using Baire category, in both the proof of A.1 and A.2, was triggered by Remark 2.27 in ([MS]) and the Category Lemma in ([To]). In fact, the commutative case A.1 below is essentially contained in ([To]). We have included the complete proof, with a different treatment of the “density”, for the reader’s convenience.

A.1. Lemma. *Let (X, μ) be a standard non atomic probability space and $\{\theta_n\}_{n \geq 1} \in \text{Aut}(X, \mu)$ a sequence of properly outer m.p. automorphisms of (X, μ) . Denote by $\mathcal{V} \subset \text{Aut}(X, \mu)$ the set of all $\theta \in \text{Aut}(X, \mu)$ with the property that $\theta_{i_0} \prod_{j=1}^n (\theta \theta_{i_{2j-1}} \theta^{-1} \theta_{i_{2j}})$ is properly outer, $\forall n \geq 1$, $i_1, i_2, \dots, i_{2n-1} \in \{1, 2, 3, \dots\}$ and $i_0, i_{2n} \in \{0, 1, 2, \dots\}$, where $\theta_0 = \text{id}_X$. Then $\mathcal{V}$ is a G_δ dense subset of $\text{Aut}(X, \mu)$. In particular, $\mathcal{V} \neq \emptyset$.*

Proof. Let $A = L^\infty(X, \mu)$, $\tau = \int \cdot d\mu$. Denote by $\mathcal{F}$ the set of all finite partitions of the identity $\{p_i\}_i \subset \mathcal{P}(A)$. If $\rho \in \text{Aut}(X, \mu)$ then we still denote by ρ the automorphism it implements on (A, τ) . As usual, $\text{Aut}(A, \tau)$ is endowed with the topology given by pointwise $\|\cdot\|_2$ convergence, with respect to which it is metrizable and complete.

For each $\rho \in \text{Aut}(A, \tau)$ denote $k(\rho) = \inf\{\|\sum_i \rho(p_i) p_i\|_2 \mid \{p_i\}_i \in \mathcal{F}\}$. Note that ρ is properly outer iff $k(\rho) = 0$. Also, if $\mathcal{D}_n$ denotes the set of $\rho \in \text{Aut}(A, \tau)$ with $k(\rho) < 1/n$ then $\mathcal{D}_n$ is clearly open in $\text{Aut}(A, \tau)$. Given an n -tuple $(\theta_1, \dots, \theta_n) \subset \text{Aut}(A, \tau)$, we denote by $\mathcal{V}_n = \mathcal{V}(\theta_1, \dots, \theta_n)$ the set of all $\rho \in \text{Aut}(A, \tau)$ with $\theta_{i_0} \prod_{j=1}^l (\rho \theta_{i_{2j-1}} \rho^{-1} \theta_{i_{2j}}) \in \mathcal{D}_n$, for all $1 \leq l \leq n$ and all choices $i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}$, $i_0, i_{2l} \in \{0, 1, 2, \dots, n\}$.

It is immediate to see that $\mathcal{V}_n$ is open in $\text{Aut}(A, \tau)$ and that $\cap_n \mathcal{V}_n = \mathcal{V}$. We have to prove that each $\mathcal{V}_n$ is also dense in $\text{Aut}(A, \tau)$, i.e., that any fixed $\rho' \in \text{Aut}(A, \tau)$ can be approximated arbitrarily well (in the point norm- $\|\cdot\|_2$ convergence on A) by some $\rho \in \mathcal{V}_n$.

By replacing if necessary $\{\theta_j\}_j$ by the properly outer automorphisms $\{\theta_j\}_j \cup \{\rho' \theta_k \rho'^{-1}\}_k$, it follows that in order to prove the density of $\mathcal{V}_n$ it is sufficient to prove that id_A is in the closure of $\mathcal{V}_n = \mathcal{V}(\theta_1, \dots, \theta_n)$, for any n -tuple of properly outer automorphisms.

To this end we will use the ultrapower II_1 factor R^ω ([McD]) as framework. Thus, we choose ω a free ultrafilter on $\mathbb{N}$ and let $R^\omega = \ell^\infty(\mathbb{N}, R)/\mathcal{I}_\omega$, where $\mathcal{I}_\omega$ is the ideal associated with the trace $\tau_\omega((x_n)_n) = \lim_{n \rightarrow \omega} \tau(x_n)$, i.e. $\mathcal{I}_\omega = \{x = (x_n)_n \mid \tau_\omega(x^* x) = 0\}$.

We regard the abelian von Neumann algebra (A, τ) as a Cartan subalgebra of R . By ([Dy]) given any $\rho \in \text{Aut}(A, \tau)$, any finite set $F \subset A$ and $\varepsilon > 0$, there exists $v \in \mathcal{N}_R(A)$ such that $\|\rho(a) - \text{Ad}(v)(a)\|_2 \leq \varepsilon$, $\forall a \in F$. Thus, there exist $u_n \in \mathcal{N}_R(A)$ such that $u = (u_n)_n \in R^\omega$ satisfies $uau^* = \rho(a)$, $\forall a \in A \subset A^\omega \subset R^\omega$. Note that ρ is properly outer iff $E_{A' \cap R^\omega}(u) = 0$.

Write $A = \overline{\cup_m D_m}^w$, for some increasing sequence of finite dimensional subalgebras of A .

Let $\mathcal{N}_m \stackrel{\text{def}}{=} D'_m \cap \mathcal{N}_R(A)$, i.e., the part of the normalizer of A in R that leaves D_m pointwise fixed. To prove that id_A is in the closure of $\mathcal{V}_n$, it is sufficient to prove:

(A.1.1). Let $U_0 = 1, U_1, U_2, \dots, U_n \in \mathcal{N}_{R^\omega}(A)$ with $E_{A' \cap R^\omega}(U_i) = 0, \forall i \neq 0$. For all $m \geq 1$, there exists $u \in \mathcal{N}_m$ such that $E_{A' \cap R^\omega}(u_{i_0} \prod_{j=1}^l (u U_{i_{2j-1}} u^* U_{i_{2j}})) = 0$, for all $1 \leq l \leq n$ and all choices $i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}, i_0, i_{2l} \in \{0, 1, 2, \dots, n\}$.

We construct u by a maximality argument, “patching together” partial isometries in $\mathcal{G}_m \stackrel{\text{def}}{=} \{vp \mid v \in \mathcal{N}_m, p \in \mathcal{P}(A)\} = D'_m \cap \mathcal{GN}_R(A)$.

Thus, for each $v \in \mathcal{G}_m$ with $vv^* = v^*v$ and each $1 \leq k \leq 2n$ we denote $\mathcal{V}_k^v$ the set of all products of the form $\prod_{j=0}^{k-1} (U_{i_j} v^{\alpha_j}) U_{i_k}$, where $i_j \in \{1, 2, \dots, n\}, \forall 1 \leq j \leq k-1, i_0, i_k \in \{0, 1, \dots, n\}, \alpha_j \in \{\pm 1\}, \alpha_1 \neq \alpha_2 \neq \dots \neq \alpha_k$. We also put $\mathcal{V}_0^v = \{U_i \mid 1 \leq i \leq n\}$. Note that if $i_0, i_{2l} \in \{0, 1, \dots, n\}, i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}$, then $U_{i_0} \prod_{j=1}^l (v U_{i_{2j-1}} v^* U_{i_{2j}}) \in \mathcal{V}_{2l}^v$. We denote

$$\mathcal{W} = \{v \in \mathcal{G}_m \mid vv^* = v^*v, E_{A' \cap R^\omega}(x) = 0, \forall x \in \mathcal{V}_k^v, 1 \leq k \leq 2n\}.$$

We endow $\mathcal{W}$ with the order given by: $v \leq v'$ if $v = v'v^*v$. $(\mathcal{W}, \leq)$ is clearly inductively ordered. Let $v_0 \in \mathcal{W}$ be a maximal element. Assume $0 \neq p = 1 - v_0 v_0^* \in \mathcal{P}(A)$. Since $E_{A'}(x) = 0$ for all $x \in \mathcal{V}_k^{v_0}, 1 \leq k \leq 2n$, it follows that $E_{A'}(pxp) = 0$ as well. Since all such elements $w = p x p$ satisfy $ww^*, w^*w \in A$ and $wAw^* = Aww^*$, by ([Dy]) it follows that

(A.1.2). $\exists 0 \neq q_1 \in \mathcal{P}(Ap)$ such that: $q_1 D_m = \mathbb{C}q_1, q_1 w q_1 = E_{A'}(w)q_1 = 0, \forall w \in \cup_{k=0}^{2n} \mathcal{V}_k^{v_0}$.

Let $v_1 \in \mathcal{N}_{q_1 R q_1}(Aq_1)$. Note that $v_1 \in \mathcal{G}_m, v_1 v_1^* = v_1^* v_1 = q_1$. Denote $u = v_0 + v_1$. We will show that if $v_1 \in \mathcal{N}_{q_1 R q_1}(Aq_1)$ is chosen appropriately then $u \in \mathcal{W}$, thus contradicting the maximality of v_0 . Write

$$x = \prod_{j=0}^{k-1} (U_{i_j} u^{\alpha_j}) U_{i_k} = \sum_{\beta} \prod_{j=0}^{k-1} (U_{i_j} v_{\beta_j}^{\alpha_j}) U_{i_k} = \sum_{\beta} y_{\beta},$$

where the sum is taken over all choices $\beta = (\beta_j)_{j=0}^{k-1} \in \{0, 1\}^k$. We will show that v_1 can be taken such that $E_{A'}(y_{\beta}) = 0, \forall \beta, \forall x \in \cup_{k=0}^{2n} \mathcal{V}_k^u$. For $\beta = (0, 0, \dots, 0)$ we have $y_{\beta} = \prod_{j=0}^{k-1} (U_{i_j} v_0^{\alpha_j}) U_{i_k} \in \mathcal{V}_k^{v_0}$, so that $E_{A'}(y_{\beta}) = 0$, by the fact that $v_0 \in \mathcal{W}$.

The k terms y_{β} corresponding to just one occurrence of v_1 (i.e. $\beta = (\beta_1, \dots, \beta_k)$ with all $\beta_i = 0$ except one), are of the form $w_0 v_1 w_1$, with $w_0, w_1 \in \cup_{j=0}^{k-1} \mathcal{V}_j^{v_0}$. Thus, each w_i satisfies $w_i w_i^*, w_i^* w_i \in A, w_i A w_i^* = A w_i w_i^*, i = 0, 1$. By shrinking q_1 recursively, we may assume $(w_0^* w_0) q_1 (w_1 w_1^*)$ is either equal to 0 or to q_1 , for all such $y_{\beta}, \forall 0 \leq k \leq 2n$. For the y_{β} for which $(w_0^* w_0) q_1 (w_1 w_1^*) = 0$ we have $y_{\beta} = 0$ and there is nothing to prove. For the y_{β} with $(w_0^* w_0) q_1 (w_1 w_1^*) = q_1$, take $u_0, u_1 \in \mathcal{N}_{R^\omega}(A)$ such that $u_0 q_1 = w_0 q_1, q_1 u_1 = q_1 w_1$. Then $E_{A'}(y_{\beta}) = E_{A'}(u_0 v_1 u_1) = u_0 E_{A'}(v_1 u_1 u_0) u_0^*$, so that $\|E_{A'}(y_{\beta})\|_1 = \|E_{A'}(v_1 u_1 u_0)\|_1$. Shrinking q_1 recursively again, all conditions so far are still satisfied while we can assume $q_1 u_1 u_0 q_1$ is either 0 or an element in $A' \cap R^\omega$, for all u_0, u_1 , coming from all w_0, w_1 arising as above. Thus, if we take $v_1 \in \mathcal{N}_{q_1 R q_1}(Aq_1)$ to be properly outer, then in both cases

$E_{A'}(v_1 u_1 u_0) = 0$, $\forall u_0, u_1$. We have thus shown that q_1, v_1 can be chosen so that for all $\beta = (\beta_1, \dots, \beta_k)$ having just one occurrence of 1 we have $E_{A'}(y_\beta) = 0$ as well.

Finally, the y_β with at least two occurrences of v_1 can be written as $y_\beta = x_0(v_1^\alpha w v_1^{\alpha'})x_1$, with $w \in \mathcal{V}_l^{v_0}$, for some $0 \leq l \leq k-2$, $\alpha, \alpha' \in \{\pm 1\}$ and x_i partial isometries. Thus

$$\|E_{A'}(y_\beta)\|_1 \leq \|y_\beta\|_1 \leq \|v_1^\alpha w v_1^{\alpha'}\|_1 = \|q_1 w q_1\|_1 = \|E_{A'}(w)q_1\|_1 = 0$$

the latter equality by (A.1.2).

Altogether, $E_{A'}(x) = 0$, $\forall x \in \cup_{k=0}^{2n} \mathcal{V}_k^u$, showing that $u \in \mathcal{W}$. But this contradicts the maximality of v_0 . Thus v_0 must be a unitary element, finishing the proof that $\mathcal{V}_n$ is dense in $\text{Aut}(A, \tau)$ and thus the proof of the statement. $\square$

A.2. Lemma. *Let $\{\theta_n\}_{n \geq 1}$ be a sequence of properly outer automorphisms of the hyperfinite II_1 factor R . Denote by $\mathcal{V}$ the set of all automorphisms θ of R such that any automorphism of R of the form $\theta_{i_0} \prod_{j=1}^n (\theta \theta_{i_{2j-1}} \theta^{-1} \theta_{i_{2j}})$ is outer, $\forall n \geq 1$, $i_1, i_2, \dots, i_{2n-1} \in \{1, 2, 3, \dots\}$ and $i_0, i_{2n} \in \{0, 1, 2, \dots\}$, where $\theta_0 = \text{id}_R$. Then $\mathcal{V}$ is a G_δ dense subset of $\text{Aut}(R)$. In particular, $\mathcal{V} \neq \emptyset$.*

Proof. Let $\{u_n\}_n$ be a sequence of unitary elements in R , dense in $\mathcal{U}(R)$ in the norm $\|\cdot\|_2$ and with each element repeated infinitely many times. For each $x \in R$ and $\rho \in \text{Aut}(R)$ denote by $k(\rho, x)$ the unique element of minimal norm $\|\cdot\|_2$ in $K(\rho, x) = \overline{\text{co}}^w \{\rho(v)xv^* \mid v \in \mathcal{U}(R)\}$. Let $\mathcal{D}_n$ be the set of automorphisms ρ of R with the property that $\|k(\rho, u_i)\|_2 < 1/n$, $\forall 1 \leq i \leq n$.

We claim that $\mathcal{D}_n$ is open in $\text{Aut}(R)$. To see this, let $\rho \in \mathcal{D}_n$ and for each $1 \leq i \leq n$ choose $v_1^i, v_2^i, \dots, v_{m_i}^i \in \mathcal{U}(R)$ such that $\|m_i^{-1} \sum_{j=1}^{m_i} \rho(v_j^i) u_i v_j^{i*}\|_2 < 1/n$. If $\delta > 0$ is sufficiently small, then any $\rho' \in \text{Aut}(R)$ satisfying $\|\rho'(v_j^i) - \rho(v_j^i)\|_2 < \delta$, $\forall 1 \leq i \leq n$, $\forall 1 \leq j \leq m_i$, will satisfy $\|m_i^{-1} \sum_{j=1}^{m_i} \rho'(v_j^i) u_i v_j^{i*}\|_2 < 1/n$, implying that $\|k(\rho', u_i)\|_2 < 1/n$, $\forall 1 \leq i \leq n$, thus $\rho' \in \mathcal{D}_n$.

Denote by $\mathcal{V}_n = \mathcal{V}_n(\theta_1, \dots, \theta_n)$ the set of all $\rho \in \text{Aut}(R)$ with the property that $\sigma_{i_0} \prod_{j=1}^l (\rho \theta_{i_{2j-1}} \rho^{-1} \theta_{i_{2j}}) \in \mathcal{D}_n$, for all $1 \leq l \leq n$ and all choices $i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}$, $i_0, i_{2l} \in \{0, 1, 2, \dots, n\}$. Since $\mathcal{D}_n$ is open and

$$\text{Aut}(R) \ni \rho \mapsto \theta_{i_0} \prod_{j=1}^l (\rho \theta_{i_{2j-1}} \rho^{-1} \theta_{i_{2j}})$$

is a continuous map, for each m and i_j as before, it follows that $\mathcal{V}_n$ is open in $\text{Aut}(R)$.

It is immediate to see that $\cap_n \mathcal{V}_n = \mathcal{V}$. Thus, in order to show that $\mathcal{V}$ is a G_δ -dense subset of $\text{Aut}(R)$ we have to prove that each $\mathcal{V}_n$ is dense in $\text{Aut}(R)$. Moreover, arguing as in the proof of A.1 we see that, by replacing if necessary $\{\theta_n\}_{n \geq 1}$ with the sequence $\{\theta'_n\}_n = \{\theta_n\}_n \cup \{\rho \theta_m \rho^{-1}\}_m$, it is enough to show that id_R is in the closure of $\mathcal{V}_n$. We will prove that in fact id_R is in the closure of $\mathcal{V}_n \cap \text{Int}(R)$, i.e., given any finite dimensional subfactor $R_0 \subset R$, there exists $u \in \mathcal{U}(R'_0 \cap R)$ such that $\text{Ad}(u) \in \mathcal{V}_n$. In turn, this will be an easy consequence of Theorem 2.1 in [P7].

To make the ideas more transparent, let us consider first the case when $\{\theta_n \mid n \geq 0\}$ is an enumeration of the automorphisms of a free cocycle action of a countable group Γ

on R , $\theta : \Gamma \rightarrow \text{Aut}(R)$. Let $M = R \rtimes \Gamma$, $U_0 = 1, U_1, U_2, \dots \in \mathcal{U}(M)$ be the canonical unitaries implementing $\theta_0, \theta_1, \dots$. Since the action is free (i.e. θ_g non-inner $\forall g \neq e$), we have $R' \cap M = \mathbb{C}$. Fix a free ultrafilter ω on $\mathbb{N}$. We view $R \subset M$ as subalgebras of constant sequences in the ultrapower II_1 factor M^ω .

Since $(R'_0 \cap R)' \cap M = R_0$, by Theorem 2.1 in [P7] there exists a unitary element $V \in (R'_0 \cap R)^\omega \subset M^\omega$ such that $VRV^* \vee M \simeq R *_R M$. In particular, if $w \in \mathcal{U}(R'_0 \cap R)$ is a Haar unitary and we put $U = VwV^* \in VRV^*$, then for any choice of $1 \leq l \leq n$, $i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}$, $i_0, i_{2l} \in \{0, 1, 2, \dots, n\}$ and $1 \leq r \leq n$ the unitary elements

$$(A.2.1) \quad \text{Ad}(U_{i_0} \Pi_{j=1}^l ((UU_{i_{2j-1}} U^{-1}) U_{i_{2j}})) (U^k) u_r U^{-k}, k = 1, 2, \dots$$

are mutually orthogonal with respect to the scalar product given by the trace. To see this, we need to show that $\tau(\text{Ad}(U_{i_0} \Pi_{j=1}^l ((UU_{i_{2j-1}} U^{-1}) U_{i_{2j}})) (U^k) u_r U^{-k}) = 0$, $\forall k \neq 0$. This amounts to $\tau(U^{-k} \Pi_{j=1}^l (UU_{i_{2j-1}} U^{-1} U_{i_{2j}}) U^k \Pi_{j=1}^l (U_{i_{2l-2j}}^{-1} U U_{i_{2l-2j+1}}^{-1} U^{-1}) u_r) = 0$, which does indeed hold true, because after some appropriate word-reduction we are left with a word of alternating “letters” $U^k \in VRV^* \ominus R_0$, $\forall k \neq 0$, and $U_{i_j}, U_{i_j}^* \in M \ominus R \subset M \ominus R_0$, for all $i_j \neq 0$.

By the orthogonality of the elements in (A.2.1), it follows that for large enough N we have

$$(A.2.2) \quad \|N^{-1} \sum_{k=1}^N \text{Ad}(U_{i_0} \Pi_{j=1}^l ((UU_{i_{2j-1}} U^{-1}) U_{i_{2j}})) (U^k) u_r U^{-k}\|_2 < 1/n,$$

for all choices of $1 \leq l \leq n$, $i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}$, $i_0, i_{2l} \in \{0, 1, 2, \dots, n\}$ and $1 \leq r \leq n$. Writing U as a sequence of unitary elements in $R'_0 \cap R$, $U = (v_m)_m$, it follows that for m large enough $u = v_m$ satisfies

$$(A.2.3) \quad \|N^{-1} \sum_{k=1}^N \theta_{i_0} \Pi_{j=1}^l (\text{Ad}(u) \theta_{i_{2j-1}} \text{Ad}(u^*) \theta_{i_{2j}}) (u^k) u_r u^{-k}\|_2 < 1/n,$$

for all $1 \leq l \leq n$, $i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}$, $i_0, i_{2l} \in \{0, 1, 2, \dots, n\}$, $1 \leq r \leq n$. Thus $\text{Ad}(u) \in \mathcal{V}_n$, finishing the proof of this particular case.

Now, in the general case we can take $\theta_0 = \text{id}_R, \theta_1, \theta_2, \dots$ to be a lifting in $\text{Aut}(R)$ of an injective group morphism $\Gamma \rightarrow \text{Out}(R)$, with Γ generated by n elements. Notice that the automorphisms $\theta_j \otimes \theta_j^{op}$ on $R \overline{\otimes} R$ implement a cocycle action $\tilde{\theta} : \Gamma \rightarrow \text{Aut}(R \overline{\otimes} R^{op})$ (see e.g. Section 3 in [P10]), so we can consider the crossed product factor $\tilde{M} = R \overline{\otimes} R^{op} \rtimes \Gamma$. Denote by $U_n \in \tilde{M}$ the canonical unitaries implementing $\theta_n \otimes \theta_n^{op}$. By [P10] again, we can view $R \overline{\otimes} R \subset \tilde{M}$ as the symmetric enveloping inclusion associated with a “diagonal subfactor” $N \subset R \simeq \mathbb{M}_{n+1}(N)$, with the embedding of N given by $x \oplus \theta_1(x) \dots \oplus \theta_n(x)$. Moreover, the associated Jones tower $N \subset R \subset N_1 \subset \dots \nearrow N_\infty$ can be viewed as a sequence of subalgebras of $\tilde{M}$, making a non-degenerate commuting square:

$$\begin{array}{ccc} R \overline{\otimes} R^{op} & \subset & \tilde{M} \\ \cup & & \cup \\ R \vee R' \cap N_\infty & \subset & N_\infty \end{array}$$

As before, we view $\tilde{M}$ as the algebra of constant sequences in $\tilde{M}^\omega$. Since $(R'_0 \cap R)' \cap N_\infty = R_0 \vee R' \cap N_\infty$ and each $R' \cap N_k$ is finite dimensional, we can apply Theorem 2.1 in [P7] to get a unitary element $V \in (R'_0 \cap R)^\omega \subset N_\infty^\omega$ such that $VN_\infty V^* \vee N_\infty \simeq N_\infty *_{R_0 \vee R' \cap N_\infty} N_\infty$. By the above commuting square, we then also have $V\tilde{M}V^* \vee \tilde{M} \simeq \tilde{M} *_{R_0 \vee R^{op}} \tilde{M}$.

Like before, take $w \in \mathcal{U}(R'_0 \cap R)$ to be a Haar unitary and denote $U = VwV^* \in VRV^*$. For any choice of $1 \leq l \leq n$, $i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}$, $i_0, i_{2l} \in \{0, 1, 2, \dots, n\}$ and $1 \leq r \leq n$, the unitary elements

$$(A.2.1') \quad \text{Ad}(U_{i_0} \Pi_{j=1}^l ((UU_{i_{2j-1}} U^{-1}) U_{i_{2j}})) (U^k) u_r U^{-k}, k = 1, 2, \dots$$

are then mutually orthogonal with respect to the scalar product given by the trace. Indeed, since $U^k \in VRV^* \ominus R_0 \subset VN_\infty V^* \ominus R_0 \vee N' \cap N_\infty$, $\forall k \neq 0$, and $U_{i_j}, U_{i_j}^* \in \tilde{M} \ominus R \vee R^{op}$, $\forall i_j \neq 0$, it follows that $\tau(\text{Ad}(U_{i_0} \Pi_{j=1}^l ((UU_{i_{2j-1}} U^{-1}) U_{i_{2j}})) (U^k) u_r U^{-k}) = 0$, for $k \neq 0$, showing the orthogonality in (A.2.1').

Now, by the orthogonality of the elements in (A.2.1'), it follows that for large enough N we have

$$(A.2.2') \quad \|N^{-1} \sum_{k=1}^N \text{Ad}(U_{i_0} \Pi_{j=1}^l ((UU_{i_{2j-1}} U^{-1}) U_{i_{2j}})) (U^k) u_r U^{-k}\|_2 < 1/n,$$

for all choices of $1 \leq l \leq n$, $i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}$, $i_0, i_{2l} \in \{0, 1, 2, \dots, n\}$ and $1 \leq r \leq n$. Writing U as a sequence of unitary elements in $R'_0 \cap R$, $U = (v_m)_m$, it follows again that for large enough m the unitary element $u = v_m$ satisfies

$$(A.2.3') \quad \|N^{-1} \sum_{k=1}^N \theta_{i_0} \Pi_{j=1}^l (\text{Ad}(u) \theta_{i_{2j-1}} \text{Ad}(u^*) \theta_{i_{2j}}) (u^k) u_r u^{-k}\|_2 < 1/n,$$

for all $1 \leq l \leq n$, $i_1, i_2, \dots, i_{2l-1} \in \{1, 2, \dots, n\}$, $i_0, i_{2l} \in \{0, 1, 2, \dots, n\}$, $1 \leq r \leq n$. Thus $\text{Ad}(u) \in \mathcal{V}_n$. □

A.3. Remarks. The proofs of both Theorems A.1 and A.2 can of course be carried out without using the hyperfinite II_1 factor R and its ultrapower R^ω as framework, working exclusively in the space $\text{Aut}(X, \mu)$, respectively $\text{Aut}(R)$. But while it is straightforward to re-write the proof of A.1 this way, the proof of A.2 then becomes much more tedious, as one can no longer use results from ([P7]). Instead, one has to go through a similar maximality argument as in the proof of A.1 but with the non-commutativity requiring some complicated estimates, similar to (pp 189-192 in [P7]).

When written in the “ $\text{Aut}(X, \mu)$ framework”, a suitable adaptation of the proof of A.1 shows the following result:

A.3.1°. Let $(X, \mathcal{X}, \nu)$ be a standard probability space and denote by $\mathcal{A}$ the set of all measurable isomorphisms ρ of X into X such that $\nu \circ \rho$ is non-singular with respect to ν . Let μ be another measure (not necessarily finite) on $(X, \mathcal{X})$, equivalent to ν . Denote by $\text{Aut}(X, \mu)$ the μ -preserving automorphisms in $\mathcal{A}$. If $\{\theta_n\}_{n \geq 1} \in \mathcal{A}$ are properly outer then the set $\mathcal{V} \subset \text{Aut}(X, \mu)$ of all $\theta \in \text{Aut}(X, \mu)$ with the property that $\theta_{i_0} \Pi_{j=1}^n (\theta \theta_{i_{2j-1}} \theta^{-1} \theta_{i_{2j}})$

is properly outer, $\forall n \geq 1$, $i_1, i_2, \dots, i_{2n-1} \in \{1, 2, 3, \dots\}$ and $i_0, i_{2n} \in \{0, 1, 2, \dots\}$, where $\theta_0 = id_X$, is a G_δ dense subset of $\text{Aut}(X, \mu)$.

In turn, the proof of the non-commutative case in the “ $\text{Aut}(R)$ framework” can be adapted to show the following more general result:

A.3.2°. For each $n \geq 1$, let $\theta_n : R^\infty \rightarrow R^\infty$ be either endomorphism or anti-endomorphism of the hyperfinite II_∞ factor R^∞ such that θ_n is outer, $Tr \circ \theta_n$ is a finite multiple of the trace Tr and $\theta_n(R^\infty)' \cap R^\infty$ is atomic, $\forall n$. Denote by $\mathcal{V}$ the set of all Tr -preserving automorphisms θ of R^∞ such that any product $\theta_{i_0} \Pi_{j=1}^n (\theta \theta_{i_{2j-1}} \theta^{-1} \theta_{i_{2j}})$ is outer, $\forall n \geq 1$, $i_1, i_2, \dots, i_{2n-1} \in \{1, 2, 3, \dots\}$ and $i_0, i_{2n} \in \{0, 1, 2, \dots\}$, where $\theta_0 = id_R$. Then $\mathcal{V}$ is a G_δ dense subset of $\text{Aut}(R^\infty)$.

The case $[R^\infty : \theta_n(R^\infty)] < \infty$, $\forall n$, of A.3.2° follows easily from Theorem 2.1 in [P7], by arguing exactly as in the proof of A.2 above: The only change in that argument is the definition of the finite index subfactor $N \subset R$, which will again be a “diagonal” inclusion, but with $R \simeq N^t$ for some appropriate amplification of N and N embedded into it by taking a partition $p_0, \dots, p_n \in \mathcal{P}(R)$ and defining $N \simeq p_0 R p$, $t = \tau(p_0)^{-1}$, $N \hookrightarrow N^t = R$ by $x + \sum_{i=1}^n \theta_i(x) p_i$, $x \in p_0 R p_0$, where $\tau(p_i)/\tau(p_0) = dTr \circ \theta_i / dTr$, $\theta_i : p_0 R p_0 \rightarrow p_i R p_i$ being “corners” of the endomorphisms $\theta_i : R^\infty \rightarrow R^\infty$. When the resulting subfactor is extremal, the rest of the argument is identical, while in case it is not extremal, then one replaces the symmetric enveloping algebra of $N \subset R$ by an appropriately defined enveloping algebra $\tilde{M}$, containing N_∞ and satisfying appropriate commuting square properties.

Note that this result shows in particular that given any two subfactors of finite Jones index of the hyperfinite II_1 factor, $P \subset R$ and $Q \subset R$, there exists a subfactor $N \subset R$ having standard invariant the “free product” of the standard invariants $\mathcal{G}_{P,R}, \mathcal{G}_{Q,R}$, as considered in ([BJ]).

REFERENCES

- [BeVa] B. Bekka, A. Valette: *Group cohomology, harmonic functions and the first L^2 -Betti numbers*, Potential Analysis, **6** (1997), 313-326.
- [BJ] D. Bisch, V.F.R. Jones: *Algebras associated to intermediate subfactors*, Invent. Math. **128** (1997), 89-157.
- [Bo] F. Boca: *Completely positive maps on amalgamated product C^* -algebras*. Math. Scand., **72** (1993), no. 2, 212-222.
- [Bu] M. Burger: *Kazhdan constants for $\text{SL}(3, \mathbb{Z})$* , J. reine angew. Math., **413** (1991), 36-67.
- [Ch] M. Choda: *A continuum of non-conjugate property (T) actions of $\text{SL}(n, \mathbb{Z})$ on the hyperfinite II_1 factor*, Math. Japon., **30** (1985), 133-150.
- [C1] A. Connes: *Classification of injective factors*, Ann. of Math., **104** (1976), 73-115.
- [C2] A. Connes: *Outer conjugacy classes of automorphisms of factors*, Ann. Ecole Norm. Sup., **8** (1975), 383-419.
- [C3] A. Connes: *A type II_1 factor with countable fundamental group*, J. Operator Theory **4** (1980), 151-153.

- [C4] A. Connes: *Sur la classification des facteurs de type II*, C. R. Acad. Sci. Paris **281** (1975), 13-15.
- [CJ] A. Connes, V. Jones: *Property T for von Neumann algebras*. Bull. London Math. Soc., **17** (1985), no. 1, 57-62.
- [D] H. Dye: *On groups of measure preserving transformations*, II, Amer. J. Math, **85** (1963), 551-576.
- [DyR] K. Dykema, F. Rădulescu: *Compressions of free products of von Neumann algebras*. Math. Ann., **316** (2000), no. 1, 61-82.
- [FM] J. Feldman, C.C. Moore: *Ergodic equivalence relations, cohomology, and von Neumann algebras I, II*, Trans. Amer. Math. Soc. **234** (1977), 289-324, 325-359.
- [Fe] T. Fernos: *Kazhdan's Relative Property (T): Some New Examples*, math.GR/0411527.
- [Fu1] A. Furman: *Orbit equivalence rigidity*, Ann. of Math. **150** (1999), 1083-1108.
- [Fu2] A. Furman: *Outer automorphism groups of some ergodic equivalence relations*, preprint 2003.
- [G1] D. Gaboriau: *Invariants ℓ^2 de relations d'équivalence et de groupes*, Publ. Math. I.H.É.S. **95** (2002), 93-150.
- [G2] D. Gaboriau: *Examples of groups that are measure equivalent to free groups*, math.DS/0503181.
- [GP] D. Gaboriau, S. Popa: *An Uncountable Family of Non Orbit Equivalent Actions of $\mathbb{F}_n$* , to appear in Journal of the AMS, math.GR/0306011.
- [Ge1] S.L. Gefter: *On cohomologies of ergodic actions of a T-group on homogeneous spaces of a compact Lie group* (Russian), in "Operators in functional spaces and questions of function theory", Collect. Sci. Works, Kiev, 1987, pp 77-83.
- [Ge2] S.L. Gefter: *Outer automorphism group of the ergodic equivalence relation generated by translations of dense subgroups of compact group on its homogeneous space*, Publ. RIMS, Kyoto Univ. **32** (1996), 517-538.
- [GoS] O. N. Golowin, L. E. Syadowsky: *Über die Automorphismengruppen der freien Produkte*, Rec. math. Moscou, N.S. **4** (1938), 505-514.
- [H] U. Haagerup: *An example of a nonnuclear C^* -algebra, which has the metric approximation property*. Invent. Math. **50** (1979), 279-293.
- [dHV] P. de la Harpe, A. Valette: "La propriété T de Kazhdan pour les groupes localement compacts", Astérisque **175** (1989).
- [Hj] G. Hjorth: *A lemma for cost attained*, preprint UCLA 2002.
- [J] V.F.R. Jones : *Index for subfactors*, Invent. Math. **72** (1983), 1-25.
- [Jol] P. Jolissaint: *Haagerup approximation property for finite von Neumann algebras*. J. Operator Theory, **48** (2002), no. 3, 549-571.
- [Ka] E. Kaniuth: *Eberhard Der Typ der regularen Darstellung diskreter Gruppen*, Math. Ann. **182** (1969) 334-339.
- [K] D. Kazhdan: *Connection of the dual space of a group with the structure of its closed subgroups*, Funct. Anal. and its Appl., **1** (1967), 63-65.
- [Ko] H. Kosaki: *Free products of measured equivalence relations*, Journal of Functional Analysis **207** (2004), 264-299.

- [LS] R. Lyndon, P. Schupp: “Combinatorial Group Theory”, Springer-Verlag, Berlin/Heidelberg/New York, 1977.
- [McD] D. McDuff: *Central sequences and the hyperfinite factor*, Proc. London Math. Soc. **21** (1970), 443-461.
- [M] G. Margulis: *Finitely-additive invariant measures on Euclidian spaces*, Ergodic. Th. and Dynam. Sys. **2** (1982), 383-396.
- [MoS] N. Monod, Y. Shalom: *Orbit equivalence rigidity and bounded cohomology*, Preprint 2002, to appear in Ann. of Math.
- [NPSa] R. Nicoara, S. Popa, R. Sasyk: *On irrational rotation HT factors*,
- [O1] N. Ozawa: *Solid von Neumann algebras*. Preprint 2003, to Appear in Acta Math., math.OA/0302082.
- [O2] N. Ozawa: *A Kurosh type theorem for type II_1 factors*, math.OA/0401121.
- [OP] N. Ozawa, S. Popa: *Some prime factorization results for II_1 factors*, Invent. Math. **156** (2004), 223-234.
- [Pe] J. Peterson: *A 1-cohomology characterization of property (T) in von Neumann algebras*, math.OA/0409527.
- [PeP] J. Peterson, S. Popa: *On the notion of relative property (T) for inclusions of von Neumann algebras*, J. Funct. Anal., **219** (2005), 469-483.
- [P1] S. Popa: *Orthogonal pairs of $*$ -subalgebras in finite von Neumann algebras*. J. Operator Theory., (1983), no. 9, 253-268.
- [P2] S. Popa: *Correspondences*, INCREST preprint 1986, unpublished, <http://www.math.ucla.edu/~popa/p/correspondences.pdf>.
- [P3] S. Popa: *Some rigidity results for non-commutative Bernoulli shifts*, J. Fnal. Analysis **230** (2006), 273-328.
- [P4] S. Popa: *Markov traces on universal Jones algebras and subfactors of finite index*, Invent. Math. **111** (1993), 375-405.
- [P5] S. Popa: *On a class of type II_1 factors with Betti numbers invariants*. Ann. of Math **163** (2006), 809-899.
- [P6] S. Popa: *Strong rigidity of II_1 factors arising from malleable actions of w -rigid groups I, II*, Invent. Math., **165** (2006), 369-408 and 409-453.
- [P7] S. Popa: *Free independent sequences in type II_1 factors and related problems*, Asterisque **232** (1995), 187-202.
- [P8] S. Popa: *Some computations of 1-cohomology groups and construction of non orbit equivalent actions*, Journal of the Inst. of Math. Jussieu **5** (2006), 309-332.
- [P9] S. Popa: *A unique decomposition result for HT factors with torsion free core*, J. Fnal. Analysis **242** (2007), 519-525 (math.OA/0401138).
- [P10] S. Popa: *Some properties of the symmetric enveloping algebras with applications to amenability and property T*, Documenta Mathematica, **4** (1999), 665-744.
- [S] Y. Shalom: *Measurable group theory*, in “Proceedings of the European Congress of Mathematicos” (Stockholm 2004), European Math Soc, Zurich 2005, 391-424.
- [Sh] D. Shlyakhtenko: *On the classification of Full Factors of Type III*, Trans. AMS **356** (2004) 4143-4159.

- [T] E. Thoma: *Eine Charakterisierung diskreter Gruppen vom Typ I*, Invent. Math. 6 **1968** 190-196.
- [To] A. Tornquist: *Orbit equivalence and actions of $\mathbb{F}_n$* , UCLA preprint 2004.
- [U1] Ueda: *Amalgamated free product over a Cartan subalgebra*, Pacific J. Math. **191** (1999), 359-392.
- [U2] Ueda: *Notes on treeability and cost in operator algebra framework*, preprint 2005, math.OA/0504262.
- [Va] A. Valette: *Group pairs with relative property (T) from arithmetic lattices*, preprint 2004 (preliminary version 2001).
- [V1] D. Voiculescu: *Symmetries of some reduced free product C^* algebras*, in “Operator algebras and their connections with topology and ergodic theory”, Springer Lecture Notes in Math **1132** (1985), pp. 566-588.
- [V2] D. Voiculescu: *The analogues of entropy and of Fisher’s information theory in free probability II*, Invent. Math. **118** (1994), 411-440 .
- [Z1] R. Zimmer: *Strong rigidity for ergodic actions of semisimple Lie groups*, Ann. of Math. **112** (1980), 511-529.
- [Z2] R. Zimmer: “Ergodic theory and semisimple groups”, Birkhäuser-Verlag, Boston 1984.

MATH.DEPT., UCLA, LA, CA 90095-155505

E-mail address: adiioana@math.ucla.edu, jpete@math.ucla.edu, popa@math.ucla.edu